

Lifetime Probability of Developing* and Dying from Cancer for 23 Sites, 2009-2011

	MALES				FEMALES			
	Developing		Dying		Developing		Dying	
	%	1 in...	%	1 in...	%	1 in...	%	1 in...
All Sites†	43.3	2	22.8	4	37.8	3	19.3	5
Brain & ONS	0.7	145	0.5	198	0.6	182	0.4	250
Breast	0.1	764	0.0	3,083	12.3	8	2.7	37
Colorectal	4.8	21	2.0	49	4.5	22	1.9	54
Esophagus	0.8	125	0.8	127	0.2	434	0.2	474
Hodgkin lymphoma	0.2	412	0.1	2,191	0.2	501	0.0	2,954
Kidney & renal pelvis	2.0	49	0.6	163	1.2	84	0.4	289
Larynx	0.6	169	0.2	489	0.1	755	0.1	1,951
Leukemia	1.7	59	1.0	98	1.2	84	0.7	138
Liver & intrahepatic bile duct	1.3	79	0.9	111	0.5	190	0.5	213
Lung & bronchus	7.4	13	6.5	15	6.2	16	5.0	20
Melanoma of skin‡	3.0	34	0.5	202	1.9	53	0.2	414
Myeloma	0.8	121	0.5	214	0.6	161	0.4	267
Non-Hodgkin lymphoma	2.4	42	0.9	115	1.9	52	0.7	144
Oral cavity & pharynx	1.6	65	0.4	258	0.7	148	0.2	551
Ovary	--	--	--	--	1.3	75	1.0	102
Pancreas	1.5	66	1.4	74	1.5	67	1.3	76
Prostate	15.0	7	2.7	38	--	--	--	--
Stomach	1.1	92	0.5	204	0.7	149	0.3	305
Testis	0.4	260	0.0	5,065	--	--	--	--
Thyroid	0.6	174	0.1	1,833	1.7	60	0.1	1,399
Urinary bladder§	3.8	26	0.9	110	1.1	88	0.3	292
Uterine cervix	--	--	--	--	0.7	154	0.2	444
Uterine corpus	--	--	--	--	2.7	37	0.6	176

* For those who are cancer free.

† All sites excludes basal cell and squamous cell skin cancers and in situ cancers except urinary bladder.

‡ Statistics are for whites.

§ Includes invasive and in situ cancer cases.

Source: Software: DevCan: Probability of Developing or Dying of Cancer Software, Version 6.7.1, National Cancer Institute, 2014.