

Leukemia: Acute Lymphocytic Overview

This overview is based on the more detailed information in our document *Leukemia-- Acute Lymphocytic*. You can get this document and other information by calling 1-800-227-2345 or visiting our website at www.cancer.org.

What is acute lymphocytic leukemia?

Leukemia is a type of cancer that starts in cells that form new blood cells. These cells are found in the soft, inner part of the bones called the *bone marrow*.

Cancer starts when cells in the body begin to grow out of control. Cells in nearly any part of the body can become cancer, and can spread to other areas of the body. To learn more about how cancers start and spread, see *What Is Cancer?*

Normal bone marrow, blood, and lymphoid tissue

To understand the different types of leukemia, it helps to know something about the blood and lymph systems. Any blood-forming cell can turn into a leukemia cell. Once that happens, the cell can grow and divide to form many new cancer cells. These cells can take over the bone marrow, spill out into the bloodstream, and spread to other organs.

Bone marrow is the soft, spongy, inner part of bones such as the skull, shoulder blades, ribs, pelvis, and bones in the spine. All of the different types of blood cells are made in the bone marrow. Bone marrow is made up of a small number of blood stem cells, blood-forming cells, fat cells, and tissues that help the blood cells grow.

Blood stem cells go through a series of changes to make new blood cells. They can develop into 1 of the 3 main types of blood cell:

• Red blood cells, which carry oxygen from the lungs to the rest of the body and carry carbon dioxide from the body to the lungs.

- Platelets, which are pieces of cells that help your blood clot
- White blood cells, which fight infection

There are different types of white blood cells. Each has its own role in fighting infection. The 3 main types of white blood cells are granulocytes, monocytes, and lymphocytes.

Lymphocytes are the main cells that make up lymphoid tissue, which is a major part of the immune system. The 2 main types of lymphocytes are called *B cells* and *T cells*. Normal T cells and B cells do different jobs within the immune system. ALL starts in early forms of B cells or T cells.

Acute lymphocytic leukemia

Acute lymphocytic leukemia (ALL) is also called *acute lymphoblastic leukemia*. The term *acute* means that the leukemia grows quickly, and if not treated, could be fatal in a few months. People with *chronic* leukemias can live years without treatment. *Lymphocytic* means it develops from early forms of lymphocytes, a type of white blood cell. This is different from acute myeloid leukemia (AML), which starts in other blood cell types found in the bone marrow. To learn about AML, see *Leukemia--Acute Myeloid*.

Other types of cancer that start in lymphocytes are known as *lymphomas* (non-Hodgkin lymphoma and Hodgkin disease). The main difference between these types of cancers is that the cancer cells in ALL are mainly in the bone marrow and blood (although they may spread to other places), while lymphomas are mainly in lymph nodes or other organs. For more information on lymphomas, see *Non-Hodgkin Lymphoma* and *Hodgkin Disease*.

Leukemia starts in the bone marrow and then moves into the blood where it can spread to other parts of the body such as the lymph nodes, spleen, liver, central nervous system (brain and spinal cord), and other organs. Other types of cancer can start in these organs and then spread to the bone marrow (or elsewhere). Those cancers are *not* leukemia. Both children and adults can get leukemia.

ALL isn't the only kind of leukemia. There are 4 main types of leukemia:

- Acute lymphocytic leukemia
- Acute myeloid leukemia
- Chronic lymphocytic leukemia
- Chronic myeloid leukemia

Knowing the exact type can help doctors better predict each patient's outlook (prognosis) and select the best treatment.

The rest of this document contains information on ALL of adults only. Chronic leukemias of adults and acute myeloid leukemia (AML) of adults are discussed in other American Cancer Society documents. For information about ALL in children, see *Childhood Leukemia*.

What are the risk factors for acute lymphocytic leukemia?

At this time we do not know the cause of most cases of acute lymphocytic leukemia (ALL). But some cases can be linked to certain *risk factors*. A risk factor is something that affects a person's chance of getting a disease. Some risk factors, like smoking, can be controlled. Others, such as a person's age, can't be changed.

But risk factors don't tell us everything. Having a risk factor, or even several risk factors, does not mean that you will get the disease. And many people who get the disease may have few or no known risk factors. Even if a person has one or more risk factors and gets cancer, it is often very hard to know what part they may have played in getting the cancer.

There are only a few known risk factors for ALL.

- Radiation exposure
- Exposure to certain chemicals
- Infection with certain viruses
- Certain inherited syndromes
- Having an identical twin who was diagnosed with ALL in the first year of life

For more information about these risk factors, see *Leukemia – Acute Lymphocytic* (*Adults*).

Can acute lymphocytic leukemia be prevented?

There is no known way to prevent leukemia at this time.

Signs and symptoms of acute lymphocytic leukemia

Certain signs and symptoms could suggest that a person might have acute lymphocytic leukemia. But tests always are needed to confirm this. And keep in mind that these symptoms are most often caused by something other than cancer.

The most common symptoms of acute lymphocytic leukemia are caused by low blood cell counts. These include:

- Feeling tired
- Feeling weak
- Feeling dizzy or lightheaded
- Shortness of breath
- Fever
- Infections that don't get better or keep coming back
- Bruising easily
- Bleeding, such as frequent or severe nosebleeds and bleeding gums

Other symptoms can include:

- Weight loss (without trying)
- Severe night sweats (that drench the sheets)
- Loss of appetite
- Fullness or swelling of the belly
- Swollen lymph nodes
- Headaches, weakness, seizures, vomiting, trouble with balance, and/or blurred vision (if leukemia cells are in the area around the brain and spinal cord)
- Trouble breathing, chest pain, and/or cough (if the leukemia is in the chest)
- Bone or joint pain

How is acute lymphocytic leukemia found?

At this time there are no special tests that can find acute lymphocytic leukemia (ALL) early. The best way to find it early is to report any possible signs or symptoms of leukemia to the doctor right away.

The doctor will ask you questions about your health (take a medical history), and do a physical exam. He or she will look for any swollen lymph nodes, any bleeding or bruising, or signs of infection. If the doctor suspects leukemia, blood tests will likely be done. If the results suggest leukemia, you will be sent to a specialist, such as a hematologist (a doctor who treats blood diseases) or an oncologist (doctor who treats cancer).

Tests to find acute lymphocytic leukemia

Most of the symptoms seen in leukemia can also be caused by other problems like infections. For this reason, your doctor will focus on finding out if you really have leukemia.

Blood cell counts

Often, the first test is a blood count. This test looks at the numbers of the different types of blood cells and at how they look under the microscope. Changes in these things can suggest leukemia.

Most people with ALL have too many white blood cells, not enough red cells, and not enough platelets. Many of the white cells will be *blasts*, a type of immature cell not normally found in the bloodstream. These cells don't work the way they should. These findings may suggest a person has leukemia, but often a sample of bone marrow cells must be looked at to be sure.

Bone marrow tests

Samples of bone marrow are obtained with tests called bone marrow aspiration and biopsy.

The samples are usually taken from the back of the hip (pelvic) bone while you lie on your stomach. The area is numbed and a needle is used to draw up a small amount of liquid bone marrow (this is the bone marrow aspiration). After that needle is removed, a slightly larger needle is moved through the bone to remove a small sliver of bone and marrow (this is the bone marrow biopsy). After the biopsy is done, pressure will be put on the site to help prevent bleeding. You will likely feel a brief, sharp pain with the aspiration, but only pressure with the biopsy.

A doctor with special training looks at the bone marrow under a microscope to see if leukemia is present and if it is, what kind of leukemia it is. Sometimes special tests that look at the cells (and even their DNA) are needed to help the doctor decide which type of leukemia a person has. You might hear some of the following terms used: cytochemistry, flow cytometry, cytogenetics, PCR, and FISH. These are complex medical and chemical tests. Your doctor can tell you which of these you might need. You can learn more about these tests in *Leukemia: Acute Lymphocytic*.

Lumbar puncture

A lumbar puncture (or spinal tap) is done to look for leukemia cells in the cerebrospinal fluid (CSF), which is the liquid around the brain and spinal cord. The doctor first numbs an area on your lower back over the spine (usually while you are curled up on your side). A small needle is then put through the skin and between the bones of the spine and into the area around the spinal cord. Then some of the liquid is taken out. The fluid is looked at to see if it has leukemia cells in it. A lumbar puncture can also be used to put drugs into the CSF to try to prevent or treat the spread of leukemia to the spinal cord and brain.

Imaging tests

Imaging tests make pictures of the inside of the body. Because leukemia does not usually form tumors, imaging tests are not always as helpful as they are for other types of cancer. For people with ALL, these tests are done more often to look for infections or other problems rather than for the leukemia itself.

X-rays: You may get a chest x-ray to check for a lung infection. The x-ray can also show swollen lymph nodes in the chest.

CT (**computed tomography**) **scans:** This is a special kind of x-ray test in which a beam moves around the body, taking pictures from different angles. A computer combines the pictures to show a slice of the body. This test can help tell if any lymph nodes or organs in your body are swollen.

A CT scanner has been described as a large donut, with a narrow table in the middle opening. You will need to lie still on the table while the scan is being done. CT scans take longer than regular x-rays, and you might feel a bit confined by the ring while the pictures are being taken.

Before the test, you may get an injection of a contrast dye, or you may be asked to drink some contrast material. This helps doctors tell normal areas from abnormal ones. A second set of pictures is then taken.

The injection can make you feel flushed or warm, in the face or elsewhere. Some people get hives (itchy bumps). A few may have more serious allergic reactions like trouble

breathing, feeling dizzy, or passing out. Be sure to tell the doctor before the scan if you have ever had allergies or a reaction to any contrast material used for x-rays. .

MRI (magnetic resonance imaging): MRI scans are very helpful in looking at the brain and spinal cord. These scans use strong magnets and radio waves instead of x-rays to make detailed pictures of the body. MRI scans take longer than CT scans. You may be placed inside a tube, which can feel confining. Newer, more open MRI machines may sometimes be another option. The MRI machine makes loud buzzing and thumping noises that you may find disturbing. Some places give you headphones to block this noise out.

Ultrasound: Ultrasound uses sound waves to make images of the organs. It can help show whether the kidneys, liver, or spleen are enlarged. It can also be used to look at lymph nodes. This is an easy test to have. For most ultrasounds you simply lie on a table and a kind of wand (transducer) is moved over the part of your body being examined.

Gallium scan and bone scan: These tests look at a slightly radioactive chemical that is put into the blood. The chemical collects in areas of cancer or infection. These areas, called "hot spots," can be picked up by a special camera. These tests are not often done in people with ALL, but they can be useful if you have bone pain that might be caused by either infection or cancer in the bones.

How is acute lymphocytic leukemia classified?

Most types of cancer are assigned a numbered stage based on the size of the tumor and how far it has spread. But leukemia does not usually form tumors, and it already involves the bone marrow and, in many cases, has also spread to other organs. The outlook for the person with acute lymphocytic leukemia (ALL) depends on other information, such as the subtype of ALL, the age of the patient, and lab test results.

The older French, American, and British (FAB) system to classify ALL was based only on the way the leukemia cells looked under the microscope after routine staining. This system has largely been replaced, as newer lab tests now allow doctors to better classify ALL.

These lab tests provide more detailed information about the subtype of ALL and the patient's outlook. They let doctors divide ALL into groups based on the type of lymphocyte (B cell or T cell) the leukemia cells come from and how mature the leukemia cells are.

The subtypes of ALL each carry a slightly different outlook, so ask your doctor what type you have and how it may affect your treatment.

Prognostic factors

As leukemia treatment has improved over the years, research has focused on why some patients have a better chance for cure than others. Certain factors, called *prognostic factors*, can give doctors a better idea of how likely standard treatment will be successful for an individual patient. These factors include

- The patient's age
- White blood cell count
- ALL subtype, certain test results
- How the ALL responds to treatment

The American Cancer Society has detailed information about these subtypes and prognostic factors in *Leukemia - Acute Lymphocytic (Adults)*.

How is acute lymphocytic leukemia treated?

Adult acute lymphocytic leukemia (ALL) is not one disease. It is really a group of diseases, and people with different subtypes vary in how they respond to treatment. Treatment choices are based on the subtype as well as on the prognostic features in "How is acute lymphocytic leukemia classified."

The treatments used most often for ALL are:

- Chemotherapy
- Targeted therapy
- Stem cell transplant

Some patients may also have surgery and radiation. The treatment of ALL often lasts for about 2 years. It can be intense, especially in the first few months of treatment, so it is important that you are treated in a center that has experience with this disease.

You might have different types of doctors involved in your care as well. The doctor in charge of your team will most likely be a hematologist, a doctor who treats blood diseases, or a medical oncologist, a doctor that treats cancer.

Thinking about taking part in a clinical trial

Clinical trials are carefully controlled research studies that are done to get a closer look at promising new treatments or procedures. Clinical trials are one way to get state-of-the art cancer treatment. In some cases they may be the only way to get access to newer

treatments. They are also the best way for doctors to learn better methods to treat cancer. Still, they are not right for everyone.

If you would like to learn more about clinical trials that might be right for you, start by asking your doctor if your clinic or hospital conducts clinical trials. You can also call our clinical trials matching service at 1-800-303-5691 for a list of studies that meet your medical needs, or see the Clinical Trials section to learn more.

Considering complementary and alternative methods

You may hear about alternative or complementary methods that your doctor hasn't mentioned to treat your cancer or relieve symptoms. These methods can include vitamins, herbs, and special diets, or other methods such as acupuncture or massage, to name a few.

Complementary methods refer to treatments that are used along with your regular medical care. Alternative treatments are used instead of a doctor's medical treatment. Although some of these methods might be helpful in relieving symptoms or helping you feel better, many have not been proven to work. Some might even be dangerous.

Be sure to talk to your cancer care team about any method you are thinking about using. They can help you learn what is known (or not known) about the method, which can help you make an informed decision. See the Complementary and Alternative Medicine section to learn more.

Help getting through cancer treatment

Your cancer care team will be your first source of information and support, but there are other resources for help when you need it. Hospital- or clinic-based support services are an important part of your care. These might include nursing or social work services, financial aid, nutritional advice, rehab, or spiritual help.

The American Cancer Society also has programs and services – including rides to treatment, lodging, support groups, and more – to help you get through treatment. Call our National Cancer Information Center at 1-800-227-2345 and speak with one of our trained specialists on call 24 hours a day, every day.

The treatment information given here is not official policy of the American Cancer Society and is not intended as medical advice to replace the expertise and judgment of your cancer care team. It is intended to help you and your family make informed decisions, together with your doctor. Your doctor may have reasons for suggesting a treatment plan different from these general treatment options. Don't hesitate to ask him or her questions about your treatment options.

Chemotherapy for acute lymphocytic leukemia

Chemotherapy (chemo) is the use of drugs to kill cancer cells. Usually the drugs are given into a vein or by mouth. Once the drugs enter the bloodstream, they spread

throughout the body. Most chemo drugs don't reach the area around the brain and spinal cord, so the drugs may also need to be put right into the cerebrospinal fluid to kill cancer cells in that area.

Chemo is the main treatment for acute lymphocytic leukemia (ALL). It generally means getting several drugs over a long period of time (often about 2 years). Treatment is separated into three phases.

Remission induction (sometimes just called induction)

The purpose of the first phase is to bring about a remission. A remission means leukemia cells are no longer found in bone marrow samples, even after the normal marrow cells return and the blood counts become normal. But this is not the same as a cure, as leukemia cells are likely to still be hiding somewhere in the body.

You will get more than one chemo drug and the doses are often high. If the ALL cells have a certain gene change, a targeted therapy drug is a part of treatment. You will most likely also have chemo put right into the spinal fluid to treat leukemia in the area around the brain and spinal cord or to keep the leukemia cells from spreading there.

Treatment in this phase can often have serious side effects, including life-threatening infections. For this reason, the doctor will watch you closely and prescribe drugs like antibiotics if needed. You may spend some or much of this time in the hospital.

If the leukemia doesn't go into remission with the first round of chemo, another round of intensive chemo will be given.

Consolidation

After the leukemia goes into remission, the next phase is often a treatment over the next few months with chemo (sometimes with targeted therapy) using many of the same drugs that were used before. This may also include chemo into the spinal fluid. This treatment phase is often easier to take than induction.

Doctors may suggest a stem cell transplant (SCT) for patients who are at a high risk of the leukemia coming back. If you will need SCT, think about having it done as part of a clinical trial at a center that has done a lot of SCT procedures. See "Stem cell transplant for acute lymphocytic leukemia" for more details

Maintenance

The last phase of treatment, called maintenance, is meant to help keep the leukemia from coming back. It uses lower doses of chemo drugs given over about 2 years. For people whose ALL cells have a certain gene change, maintenance often includes a targeted therapy drug. Chemo may also continue to be given into the spinal fluid.

Side effects of chemo

Chemo drugs kill fast-growing cells such as cancer cells, but in the process they also damage other normal cells that grow fast.

The side effects of chemo depend on the type and dose of drugs given and how long they are taken. Common side effects might include:

- Hair loss
- Mouth sores
- Higher risk of infection (from low white blood cells)
- Easy bruising or bleeding (from low blood platelets)
- Tiredness (from low red blood cells)
- Loss of appetite
- Nausea
- Vomiting
- Diarrhea

Most side effects usually go away after treatment ends. Be sure to talk to your doctor if you are having trouble with side effects because there are often ways to manage them during treatment. For example, there are drugs that can be taken along with the chemo to help prevent or reduce nausea and vomiting. Drugs called *growth factors* are sometimes given to keep your blood counts higher and reduce the chance of infection.

Chemo for ALL can damage kidneys, liver, testicles, ovaries, brain, heart, and lungs. Your doctor will watch you carefully to try to prevent much of the damage. If serious side effects happen, the drugs may have to be reduced or stopped. Be sure to tell your doctor about any problems you have. .

Targeted therapy for acute lymphocytic leukemia

In recent years, new drugs that target certain parts of cancer cells have been developed. These targeted therapies work differently from standard chemotherapy (chemo) drugs and often have different and less severe side effects.

Sometimes acute lymphocytic leukemia (ALL) cells have a certain gene change that is similar to the gene change seen in chronic myeloid leukemia (CML) cells. When they do have this gene change, some of the targeted drugs used to treat CML are useful in treating

ALL. These drugs can help more ALL patients go into a remission and may help keep the leukemia from coming back.

A common side effect of targeted therapy drugs is swelling around the eyes or in the hands or feet. Other possible side effects include diarrhea, nausea, muscle pain, extreme tiredness (fatigue), and skin rashes, as well as lower red blood cell and platelet counts at the start of treatment.

Monoclonal antibodies to treat acute lymphoblastic leukemia

Monoclonal antibodies are man-made versions of immune system proteins (antibodies) designed to attach to a certain part of a cancer cell. They can help kill the cancer cells.

Blinatumomab (BlincytoTM), which can be used to treat some kinds of acute lymphocytic leukemia, is a special kind of monoclonal antibody because it can attach to both the leukemia cell and an immune cell. This can lead to the immune cell killing the leukemia cell.

This drug is given as an infusion into a vein, and sometimes has very serious side effects.

More detailed information can be found in Leukemia: Acute Lymphocytic (in Adults).

Surgery for acute lymphocytic leukemia

Surgery plays a very small part in the treatment of acute lymphocytic leukemia (ALL). Because leukemia cells spread widely throughout the bone marrow and to many other organs, it is not possible to cure this type of cancer by surgery.

Surgery can be used to help give treatment. To lower the number of needle sticks you'll need during treatment, a plastic tube called a *venous access device* can be placed in a large vein. This tube makes it easier to put drugs into the blood stream or remove blood samples. If you get a venous access device, you will be shown how to take care of it to prevent it from getting infected.

If you will be getting chemotherapy in the fluid surrounding the brain and spinal cord (the cerebrospinal fluid or CSF), you might also have a device called an *Ommaya reservoir* placed. This device is made up of a small dome that sits under the skin of the scalp, with the tube going through a hole in the skull and into the CSF in one of the cavities of the brain. An Ommaya reservoir also lets doctors take samples of CSF for testing without doing a lumbar puncture/spinal tap. The device stays in place until treatment is done.

Radiation therapy for acute lymphocytic leukemia

Radiation therapy is the use of high energy x-rays to kill cancer cells. It is sometimes used to treat leukemia that has spread to the brain and spinal cord or to the testicles. It might also be used to reduce pain when the leukemia has spread to a bone if chemo hasn't helped

Radiation to the whole body is often done as part of a stem cell transplant (see "Stem cell transplant for acute lymphocytic leukemia"). It is also used, though rarely, in an emergency to shrink a tumor if it is pressing on the windpipe. But more often chemo is used instead.

The possible side effects of radiation depend on where it is aimed. There may be sunburn-like skin changes in the treated area. Radiation to the belly (abdomen) can sometimes cause nausea, vomiting, or diarrhea. For radiation that includes large parts of the body, the effects may include lowered blood cell counts, which can lead to extreme tiredness (called *fatigue*) and an increased risk of infection.

Stem cell transplant for acute lymphocytic leukemia

Very high doses of chemotherapy (chemo) drugs might work better to kill acute lymphocytic leukemia (ALL) cells, but they can severely damage normal bone marrow cells, which could be life-threatening. Stem cell transplants (SCT) offer a way for doctors to use high doses of chemo. Although the drugs destroy the bone marrow, transplanted stem cells can restore the bone marrow's ability to make blood.

Stem cells for a transplant come from either the blood or from the bone marrow. They can also come from the umbilical cord of a newborn baby.

These blood-forming stem cells can come from either the patient or from a donor whose tissue type closely matches the patient's type. For people with ALL, a donor (or allogeneic) transplant is most often used. The donor may be a brother or sister or – less often – a person not related to the patient. If the patient's own cells are used it is called an autologous stem cell transplant, but this is not often done for ALL because it is hard to separate normal stem cells from leukemia cells in the bone marrow or blood. That means there is the risk of returning some leukemia cells with an autologous stem cell transplant.

To learn more about stem cell transplants and their side effects, see *Stem Cell Transplant* (*Peripheral Blood, Bone Marrow, and Cord Blood Transplants*).

Palliative treatment for acute lymphoblastic leukemia

If at some point it becomes clear that further treatment is very unlikely to cure the leukemia, it may be time to focus on relieving symptoms. Treatment aimed at symptom

control is known as *palliative treatment*. One example is milder chemo to try to slow the growth of the leukemia in order to reduce symptoms.

Other palliative treatments can include treating pain with drugs or radiation and using medicines or blood transfusions to treat low blood counts and tiredness. Nausea and loss of appetite may be helped by medicines and high-calorie food supplements. Antibiotics may be used to treat infection.

These can help you feel better and may even help you live longer, but they aren't aimed at curing the cancer.

What are some questions I can ask my doctor about acute lymphocytic leukemia?

It is important to have open discussions with your doctor. Feel free to ask any question you have, no matter how small it might seem. Here are some questions you might want to ask. Be sure to add your own questions as you think of them. Nurses, social workers, and other members of the treatment team may also be able to answer many of your questions.

- Would you please write down the exact type of leukemia I have?
- What factors might affect my outlook?
- Are there other tests that need to be done before we can decide on treatment?
- Are there other doctors I need to see?
- How much experience do you have treating this type of leukemia?
- Should I get a second opinion before starting treatment? Can you suggest someone?
- What are my treatment choices?
- Which treatment do you recommend, and why?
- How soon do we need to start treatment?
- Should we think about a stem cell transplant? When?
- What risks and side effects are there to the treatments you recommend?
- What can I do to help reduce the side effects I may have from the chemo?
- How long will treatment last? What will it be like? Where will it be done?
- How will treatment affect my daily activities?
- What are the chances that my leukemia will come back once I am in remission?

- What is my outlook?
- What will we do if the treatment doesn't work or if the leukemia comes back?
- What type of follow-up will I need after treatment?

Be sure to write down any questions you have that are not on this list.

Moving on after treatment for acute lymphocytic leukemia

For some people with acute lymphocytic leukemia (ALL), treatment may get rid of the cancer. Completing treatment can be both stressful and exciting. You may be relieved to finish treatment, but find it hard not to worry about the leukemia coming back. (When leukemia comes back after treatment, it is called *recurrence*.) This is a very common concern in people who have had leukemia.

It may take a while before your recovery begins to feel real and your fears are somewhat relieved. You can learn more about what to look for and how to learn to live with the chance of cancer coming back in *Living With Uncertainty: The Fear of Cancer Recurrence*.

For other people, the leukemia may not go away completely. These people may get regular treatments with chemo, radiation, or other treatments to help keep the leukemia in check for as long as possible. Learning to live with cancer that does not go away can be hard and very stressful. It has its own type of uncertainty. See *When Cancer Doesn't Go Away* for more about this.

Follow-up care

Treatment for ALL can last for years. Even if you have finished treatment, your doctors will still want to watch you closely. It is very important to go to all of your follow-up doctor visits. During these visits, your doctors will ask questions about any problems you may have and do an exam. Often, blood tests will be checked to look for signs of leukemia or treatment side effects.

Almost any cancer treatment can have side effects. Some may last for a few weeks or months, but others can be permanent. Please tell your cancer care team about any symptoms or side effects that bother you so they can help you manage them. Use this time to ask your health care team questions and discuss any concerns you might have.

If the leukemia does come back, it usually happens during treatment or shortly after treatment ends. It is unusual for the leukemia to return if there are still no signs of the disease 5 years after treatment.

It is also important to keep health insurance. While you hope your cancer won't come back, it could happen. If it does, you don't want to have to worry about paying for treatment. Should your cancer come back, see *When Your Cancer Comes Back: Cancer Recurrence* for information about how to manage and cope with this phase of your treatment.

Seeing a new doctor

At some point after your leukemia is found and treated, you may find yourself in the office of a new doctor. It is important that you be able to give your new doctor the exact details of your diagnosis and treatment. Gathering these details soon after treatment may be easier than trying to get them at some point in the future. Make sure you have this information handy and always keep copies for yourself:

- A copy of your pathology report from any biopsy or surgery
- If you had surgery, a copy of your operative report
- If you stayed in the hospital, a copy of the discharge summary that the doctor wrote when you were sent home
- If you had radiation treatment, a copy of your treatment summary
- If you had chemotherapy or targeted therapy drugs, a list of your drugs, drug doses, and when you took them

The doctor may want copies of this information for his records, but always keep copies for yourself.

Lifestyle changes after treatment for acute lymphocytic leukemia

You can't change the fact that you have had leukemia. What you can change is how you live the rest of your life – making choices to help you stay healthy and feel as well as you can. This can be a time to look at your life in new ways. Maybe you are thinking about how to improve your health over the long term. Some people even start during cancer treatment.

Make healthier choices

For many people, a diagnosis of cancer helps them focus on their health in ways they may not have thought much about in the past. Are there things you could do that might make you healthier? Maybe you could try to eat better or get more exercise. Maybe you could cut down on alcohol, or give up tobacco. Even things like keeping your stress level under control may help. Now is a good time to think about making changes that can have

positive effects for the rest of your life. You will feel better and you will also be healthier.

You can start by working on those things that worry you most. Get help with those that are harder for you. For instance, if you are thinking about quitting smoking and need help, call us at 1-800-227-2345.

Eating better

Eating right can be hard for anyone, but it can get even tougher during and after cancer treatment. Treatment may change your sense of taste. Nausea can be a problem. You may not feel like eating and lose weight when you don't want to. Or you may have gained weight that you can't seem to lose. All of these things can be very frustrating.

If treatment caused weight changes or eating or taste problems, do the best you can and keep in mind that these problems usually get better over time. You may find it helps to eat small portions every 2 to 3 hours until you feel better. You may also want to ask your cancer team about seeing a dietitian, an expert in nutrition who can give you ideas on how to deal with these treatment side effects.

One of the best things you can do after treatment is to put healthy eating habits into place. You may be surprised at the long-term benefits of some simple changes. Getting to and staying at a healthy weight, eating a healthy diet, and limiting your alcohol intake may lower your risk for a number of types of cancer, as well as having many other health benefits.

Fatigue and exercise

Feeling very tired (fatigue) is a very common problem during and after cancer treatment. This is not a normal type of tiredness but a "bone-weary" exhaustion that doesn't get better with rest. For some people, fatigue lasts a long time after treatment and can keep them from staying active. But exercise can actually help reduce fatigue and the sense of depression that sometimes comes with feeling so tired.

If you were very ill or weren't able to do much during treatment, it is normal that your fitness, staying power, and muscle strength declined. You need to find an exercise plan that fits your own needs. Talk with your health care team before starting anything. Get their input about your exercise plans. Then, try to find an exercise buddy so you're not doing it alone. Having family or friends involved when starting a new exercise program can give you that extra boost of support to keep you going when the push just isn't there.

If you are very tired, though, you will need to balance activity with rest. It is OK to rest when you need to. To learn more about fatigue, see *Fatigue in People With Cancer*.

Exercise can improve your physical and emotional health.

- It improves your cardiovascular (heart and circulation) fitness.
- It can help you stay at a healthy weight.
- It makes your muscles stronger.
- It reduces fatigue.
- It can help lower anxiety and depression.
- It can make you feel happier.
- It can help you feel better about yourself.

Long term, we know that getting regular physical activity plays a role in helping to lower the risk of some cancers, as well as having other health benefits.

Can I lower my risk of the leukemia progressing or coming back?

Most people want to know if there are certain lifestyle changes they can make to reduce their risk of cancer growing or coming back. Unfortunately, for most cancers there is little solid evidence to guide people. This doesn't mean that nothing will help – it's just that for the most part this is something that hasn't been well studied. Most studies have looked at lifestyle changes as ways of preventing cancer in the first place, not slowing it down or keeping it from coming back.

At this time, not enough is known about acute lymphocytic leukemia to say for sure if there are things you can do that will be helpful. Healthy behaviors such as not smoking, eating well, and staying at a healthy weight may help, but no one knows for sure. But we do know that these types of changes can have positive effects on your health that can extend beyond your risk of leukemia or other cancers.

How does having acute lymphocytic leukemia affect your emotional health?

During and after treatment, you may be surprised by the flood of emotions you go through. This happens to a lot of people. You may find that you think about the effect of your cancer on things like your family, friends, and career. You may take a new look at your relationships with those around you. Money may be a concern as the medical bills pile up. Unexpected issues may also cause concern – for instance, as you get better and need fewer doctor visits, you will see your health care team less often. This can be hard for some people.

This is a good time to look for emotional and social support. You need people you can turn to. Support can come in many forms: family, friends, cancer support groups, church or spiritual groups, online support communities, or private counselors.

The cancer journey can feel very lonely. You don't need to go it alone. Your friends and family may feel shut out if you decide not to include them. Let them in – and let in anyone else who you feel may help. If you aren't sure who can help, call your American Cancer Society at 1-800-227-2345 and we can put you in touch with a group or resource that may work for you.

If treatment for acute lymphocytic leukemia stops working

When a person has had many different treatments and the cancer has not been cured, over time the cancer tends to resist all treatment. At this time you may have to weigh the possible benefits of a new treatment against the downsides, like treatment side effects.

This is likely to be the hardest time in your battle with cancer – when you have tried everything within reason and it's just not working anymore. Your doctor may offer you new treatment, but you will need to talk about whether the treatment is likely to improve your health or change your outlook for survival.

No matter what you decide to do, it is important for you to feel as good as possible. Make sure you are asking for and getting treatment for pain, nausea, or any other problems you may have. This type of treatment is called *palliative treatment*. It helps relieve symptoms but is not meant to cure the cancer.

At some point you may want to think about hospice care. Most of the time hospice care is given at home. Your cancer may be causing symptoms or problems that need to be treated. Hospice focuses on your comfort. You should know that while getting hospice care often means the end of treatments such as chemo and radiation, it doesn't mean you can't have treatment for the problems caused by your cancer or other health issues. It just means that the purpose of your care is to help you live life as fully as possible and to feel as well as you can. You can learn more about this in *Hospice Care*.

Staying hopeful is important, too. Your hope for a cure may not be as bright, but there is still hope for good times with family and friends – times that are filled with happiness and meaning. Pausing at this time in your cancer treatment gives you a chance to focus on the most important things in your life. Now is the time to do some things you've always wanted to do and to stop doing the things you no longer want to do. Though the cancer may be beyond your control, there are still choices you can make.

What's new in acute lymphocytic leukemia research?

Researchers at many medical centers, hospitals, and other institutions are looking at the causes, diagnosis, and treatment of acute lymphocytic leukemia (ALL).

Finding "hidden" disease (minimal residual disease)

Progress has been made in finding leukemia cells after treatment when there may be so few leukemia cells that they cannot be found by routine bone marrow tests. A test called polymerase chain reaction (PCR) can find one cancer cell among many thousands of normal cells. This is helpful in seeing how well the chemo has destroyed the leukemia cells and if a relapse is likely. Doctors are now trying to figure out whether patients with minimal residual disease will be helped by further or more intense treatment.

Better chemotherapy

Studies are going on to find the best combinations of chemotherapy (chemo) drugs (those that work the best and cause the fewest side effects) and to figure out which patients will be helped the most from different types of treatment. Sometimes chemo does not work very well because the leukemia cells become resistant to it. Researchers are now looking at ways to prevent or reverse this resistance by using other drugs along with chemo. New chemo drugs are also being developed and tested.

Stem cell transplants

Studies are also being done to improve the stem cell transplant process and to predict which patients are most likely to be helped by this treatment.

Monoclonal antibodies

These are man-made immune system proteins that attach to certain molecules on the surface of the leukemia cells. Some of them are already used to treat certain lymphomas. Researchers are now looking at whether they might be helpful against ALL. Early results have been good, but it is still too early to know for sure.

More information about acute lymphocytic leukemia

We have a lot more information that you might find helpful. Explore www.cancer.org or call our National Cancer Information Center_toll-free number, 1-800-227-2345. We're here to help you any time, day or night.

National organizations and Web sites*

Along with the American Cancer Society, other sources of information and support include:

Acute lymphocytic leukemia

Leukemia & Lymphoma Society

Toll-free number: 1-800-955-4572

Website: www.lls.org

National Cancer Institute

Toll-free number: 1-800-4-CANCER (1-800-422-6237)

Website: www.cancer.gov

National Coalition for Cancer Survivorship

Toll-free number: 1-888-650-9127

1-877-NCCS-YES (622-7937) for some publications and Cancer Survivor Toolbox®

orders

Website: www.canceradvocacy.org

Bone marrow and peripheral blood stem cell transplants

National Bone Marrow Transplant Link (nbmtLINK)

Toll-free number: 1-800-LINK-BMT (1-800-546-5268)

Website: www.nbmtlink.org

Be the Match (formerly National Marrow Donor Program)

Toll-free number: 1-800-MARROW-2 (1-800-627-7692)

Website: www.bethematch.org

No matter who you are, we can help. Contact us anytime, day or night, for information and support. Call us at **1-800-227-2345** or visit www.cancer.org.

Last Medical Review: 12/8/2014

Last Revised: 2/22/2016

2014 Copyright American Cancer Society

For additional assistance please contact your American Cancer Society 1-800-227-2345 or www.cancer.org

^{*}Inclusion on this list does not imply endorsement by the American Cancer Society.