



Lung Cancer (Small Cell) Overview

The information that follows is an overview of this type of cancer. It is based on the more detailed information in our document, *Lung Cancer (Small Cell)*. This document and other information can be obtained by calling 1-800-227-2345 or visiting our Web site at www.cancer.org.

What is cancer?

The body is made up of trillions of living cells. Normal body cells grow, divide, and die in an orderly way. During the early years of a person's life, normal cells divide faster to allow the person to grow. After the person becomes an adult, most cells divide only to replace worn-out, damaged, or dying cells.

Cancer begins when cells in a part of the body start to grow out of control. There are many kinds of cancer, but they all start because of this out-of-control growth of abnormal cells.

Cancer cell growth is different from normal cell growth. Instead of dying, cancer cells keep on growing and form new cancer cells. These cancer cells can grow into (invade) other tissues, something that normal cells cannot do. Being able to grow out of control and invade other tissues are what makes a cell a cancer cell.

In most cases the cancer cells form a tumor. But some cancers, like leukemia, rarely form tumors. Instead, these cancer cells are in the blood and bone marrow.

When cancer cells get into the bloodstream or lymph vessels, they can travel to other parts of the body. There they begin to grow and form new tumors that replace normal tissue. This process is called *metastasis* (muh-**tas**-tuh-sis).

No matter where a cancer may spread, it is always named for the place where it started. For instance, breast cancer that has spread to the liver is still called breast cancer, not liver cancer. Likewise, prostate cancer that has spread to the bone is called metastatic prostate cancer, not bone cancer.


Different types of cancer can behave very differently. For example, lung cancer and breast cancer are very different diseases. They grow at different rates and respond to different treatments. That is why people with cancer need treatment that is aimed at their own kind of cancer.

Not all tumors are cancerous. Tumors that aren't cancer are called *benign* (be-**nine**). Benign tumors can cause problems – they can grow very large and press on healthy organs and tissues. But they cannot grow into other tissues. Because of this, they also can't spread to other parts of the body (metastasize). These tumors are almost never life threatening.

What is small cell lung cancer?

Note: *This document covers only the small-cell type of lung cancer. The treatment for small cell lung cancer is very different from the treatment for non-small cell lung cancer. Much of the information for one type will not apply to the other type. If you don't know which type of lung cancer you have, ask your doctor so you can be sure you are looking at the right information.*

Lung cancer is a cancer starts in the lungs. To understand lung cancer, it helps to know something about the lungs and how they work.



The lungs

The lungs are 2 sponge-like organs found in the chest. When you breathe in, air enters through your mouth and nose and goes into your lungs through the windpipe (*trachea*). The trachea divides into tubes called the *bronchi*, which divide into smaller branches called the *bronchioles*. At the end of the bronchioles are tiny air sacs known as *alveoli*. Many tiny blood vessels run through the alveoli. They absorb oxygen from the air you breathe in and pass carbon dioxide from the body into the alveoli to be breathed out when you exhale. Taking in oxygen and getting rid of carbon dioxide are your lungs' main functions.

The lining around the lungs, called the *pleura*, helps to protect the lungs and allows them to move during breathing.

Below the lungs, a muscle called the *diaphragm* separates the chest from the belly (abdomen). When you breathe, the diaphragm moves up and down, forcing air in and out of the lungs.

Start and spread of lung cancer

Most lung cancer starts in the cells lining the lung airways.

Lung cancers are thought to start as areas of pre-cancer changes in the lung. These changes are not a mass or tumor. They can't be seen on an x-ray and they don't cause symptoms.

Over time, these changes in the cells may go on to become true cancer. The cancer makes chemicals that cause new blood vessels to form nearby. These new blood vessels feed the cancer cells and allow a tumor to form. In time, the tumor becomes large enough to show up on an x-ray.

At some point, lung cancer cells can break away and spread to other parts of the body in a process called metastasis. Lung cancer is often a life-threatening disease because it can spread this way before it is found.

The lymph system

One of the ways lung cancer can spread is through the lymph (pronounced "limf") system. Lymph vessels are like veins, but they carry lymph instead of blood. Lymph is a clear fluid that contains tissue waste and cells that fight infection. Lung cancer cells can enter lymph vessels and begin to grow in lymph nodes (small collections of immune cells) around the bronchi and in the space between the lungs. When lung cancer cells have reached the lymph nodes, they are more likely to have spread to other organs of the body. The extent (stage) of the cancer and decisions about treatment are based on whether or not the cancer has spread to the nearby lymph nodes. We talk about this more in the section, "After the tests: Staging."

Types of lung cancer

There are 2 main types of lung cancer and they are treated very differently.

- Small cell lung cancer (SCLC)
- Non-small cell lung cancer (NSCLC)

(If the cancer has features of both types, it is called mixed small cell/large cell cancer. This is not common.)

The information here only covers small cell lung cancer. Non-small cell lung cancer is covered in our document, *Lung Cancer (Non-Small Cell)*.

Small cell lung cancer (SCLC)

About 10% to 15% of all lung cancers are the small cell type. Other names for SCLC are *oat cell carcinoma* and *small cell undifferentiated carcinoma*.

This cancer often starts in the bronchi near the center of the chest. Although the cancer cells are small, they can divide quickly, form large tumors, and spread to lymph nodes and other organs throughout the body. This is important because it means that surgery is rarely an option and never the only treatment given. On the other hand, chemotherapy,

which can reach cancer cells throughout the body, is an important part of treatment for all small cell lung cancers, as long as a person is healthy enough to have it.

Other types of lung cancer

Along with the 2 main types of lung cancer, other tumors can also be found in the lungs. Some of these are not cancer and others are. Carcinoid tumors, for instance, are slow-growing and usually cured by surgery. We have more information about lung carcinoid tumors in our document *Lung Carcinoid Tumor*.

Keep in mind that cancer that starts in other organs (such as the breast, pancreas, kidney, or skin) can sometimes spread (metastasize) to the lungs, but these are **not** lung cancers. For instance, cancer that starts in the breast and spreads to the lungs is still breast cancer, not lung cancer. Treatment for these cancers that have spread to the lungs depends on where the cancer started.

How many people get lung cancer?

The American Cancer Society's most recent estimates for lung cancer (both small cell and non-small cell) in the United States are for 2012:

- About 226,160 new cases of lung cancer (both small cell and non-small cell)
- About 160,340 deaths from lung cancer

Lung cancer (both small cell and non-small cell) is by far the leading cause of cancer death for both men and women. More people die of lung cancer than of colon, breast, and prostate cancers combined. Lung cancer is fairly rare in people under the age of 45.

The average lifetime chance that a man will develop lung cancer is about 1 in 13. For a woman it is 1 in 16. These numbers include both smokers and non-smokers. For smokers the risk is much higher, while for non-smokers the risk is lower.

Survival statistics based on the stage (extent) of the cancer are covered in the section, "Survival rates for small cell lung cancer."

What are the risk factors for small cell lung cancer?

A risk factor is anything that affects a person's chance of getting a disease such as cancer. Different cancers have different risk factors. Some risk factors, like smoking, can be controlled. Others, like a person's age or family history, can't be changed.

But risk factors don't tell us everything. Having a risk factor, or even several risk factors, does not mean that you will get the disease. And some people who get the disease may not have had any known risk factors. Even if a person with lung cancer has a risk factor,

it is often very hard to know how much that risk factor may have contributed to the cancer. Still, having several risk factors can make you more likely to get lung cancer.

Tobacco smoke

Smoking is by far the leading risk factor for lung cancer. Tobacco smoke causes nearly 8 out of 10 cases of lung cancer. This number is probably even higher for small cell lung cancer. It is very rare for someone who has never smoked to have small cell lung cancer. The longer a person has been smoking and the more packs per day smoked, the greater the risk. If a person stops smoking before lung cancer starts, the lung tissue slowly repairs itself. Stopping smoking at any age may lower the risk of lung cancer and help you live longer.

Cigar and pipe smoking are almost as likely to cause lung cancer as is cigarette smoking. And smoking low tar or "light" cigarettes increases lung cancer risk as much as regular cigarettes. There is concern that menthol cigarettes may increase the risk even more since the menthol allows smokers to inhale more deeply.

Secondhand smoke: People who don't smoke but who breathe the smoke of others may also be at a higher risk for lung cancer. Non-smokers who live with a smoker, for instance, have about a 20% to 30% greater risk of developing lung cancer. Non-smokers exposed to tobacco smoke in the workplace are also more likely to get lung cancer. Some people seem to be more prone to the cancer-causing effect of tobacco smoke than others.

Radon

Radon is a radioactive gas made by the normal breakdown of uranium in soil and rocks. Uranium is found at higher levels in the soil in some parts of the United States. Radon can't be seen, tasted, or smelled. It can build up indoors and create a possible risk for cancer. The lung cancer risk from radon is much lower than that from tobacco smoke. But the risk is much higher among smokers than for those who don't smoke. State and local offices of the EPA (Environmental Protection Agency) can give you information on how to test for radon in the home. The document, *Radon*, is also available from the ACS.

Asbestos

Asbestos is another risk factor for lung cancer. People who work with asbestos have a higher risk of getting lung cancer. If they also smoke, the risk is greatly increased. Both smokers and non-smokers exposed to asbestos also have a greater risk of getting a type of cancer called mesothelioma, which starts in the lining of the lungs. Because it is not really lung cancer, mesothelioma is discussed in our document, *Malignant Mesothelioma*.

Although asbestos was used for many years, the government has now nearly stopped its use in the workplace and in home products. While it is still present in many buildings, it is not thought to be harmful as long as it is not released into the air. To learn more, see our document, *Asbestos*.

Other cancer-causing things in the workplace

Other things that cause cancer (carcinogens) found in some workplaces that can increase lung cancer risk include:

- Radioactive ores, such as uranium
- Inhaled chemicals or minerals like arsenic, beryllium, cadmium, vinyl chloride, nickel compounds, chromium compounds, coal products, mustard gas, and chloromethyl ethers
- Diesel exhaust

The government and industry have taken major steps in recent years to help protect workers. But the dangers are still there. If you work around any of these, you should be very careful to limit how much you are exposed.

Radiation treatment to the lungs

People who have had radiation to the chest to treat other cancers are at higher risk for lung cancer, especially if they smoke. Women who have radiation to the breast after a lumpectomy for breast cancer do not appear to have a higher risk of lung cancer.

Arsenic

High levels of arsenic in drinking water may increase the risk of lung cancer. The effect is even greater for smokers.

Personal and family history

If you have had lung cancer, you have a higher risk of getting another lung cancer. Brothers, sisters, and children of people who have had lung cancer may have a slightly higher risk themselves, especially if the family got cancer at a younger age. Researchers have found that genes do seem to play a role in some families with a strong history of lung cancer. More research is being done on this.

Certain vitamins

Two large studies have found that smokers who took beta carotene supplements actually had an increased risk of lung cancer. The results of these studies suggest that smokers should not take beta carotene supplements.

Air pollution

In cities, air pollution may slightly increase the risk of lung cancer. But the risk is still far less than that caused by smoking. Worldwide, about 5% of all deaths from lung cancer may be due to outdoor air pollution.

DNA and gene changes

During the past few years, scientists have made great progress in learning how risk factors produce certain changes in the DNA of lung cells, causing the cells to become cancer. DNA is the genetic material that carries the instructions for nearly everything our cells do.

Current research in this field is aimed at developing tests that can find lung cancers at an early stage by spotting DNA changes. But these tests are not yet ready for routine use. For now, doctors stress the importance of not using tobacco and staying away from other things that can increase your cancer risk.

Can small cell lung cancer be prevented?

Some people who get lung cancer do not have any known risk factors. Although we know how to prevent most lung cancers, at this time we don't know how to prevent all of them.

The best way to reduce your risk of lung cancer is not to smoke. You should also avoid breathing in other people's smoke. If you would like help quitting smoking, see our *Guide to Quitting Smoking* or call the American Cancer Society at 1-800-227-2345.

Radon is also a cause of lung cancer. You can lower your exposure by having your home tested and treated, if needed. For more information, see our document, *Radon*.

Protecting yourself from cancer-causing chemicals at work and elsewhere can also be helpful. When people work where these exposures are common, they should be kept as low as possible.

A good diet with lots of fruits and vegetables may also help reduce your risk of lung cancer.

How is small cell lung cancer found?

It is often hard to find lung cancer early. Most people with early lung cancer do not have any symptoms, so only a small number of lung cancers are found at an early stage. When lung cancer is found early, it is often because of tests that were being done for something else.

Does screening for lung cancer save lives?

Screening is the use of tests or exams to find a disease like cancer in people who don't have any symptoms. Because lung cancer often spreads beyond the lungs before it causes symptoms, a good screening test to find lung cancer early could save many lives.

For many years, doctors have been trying to figure out if finding lung cancer early can save lives. Before now, no lung cancer screening test had been shown to lower the risk of dying from this disease. Studies of 2 screening tests, chest x-ray and sputum cytology,

did find that these tests could find lung cancers at an early stage, but neither test helped patients live longer. For this reason, major medical groups have not recommended routine screening tests to find lung cancer early, even for people at increased risk, such as smokers.

Now, though, a different lung cancer screening test has been shown to help lower the risk of dying from this disease.

Low-dose spiral CT

A special type of CT scan, known as low-dose spiral CT (or helical CT) has shown some promise in finding early lung cancers in heavy smokers and former smokers. Spiral CT gives more detailed pictures than a chest x-ray and is better at finding small changes in the lungs. The type used to screen for lung cancer is called “low-dose” because it uses less radiation than the standard spiral CT of the chest.

The National Lung Screening Trial (NLST) is a large study that compared spiral CT scans to chest x-rays in people at high risk of lung cancer to see if these scans could help lower the risk of dying from lung cancer. People in the study were current or former heavy smokers aged 55 to 74. They got either 3 spiral CT scans or 3 chest x-rays, each a year apart. They were then followed for several years to see how many people in each group died of lung cancer.

In the study, the people who got spiral CT had a 20% lower chance of dying from lung cancer than those who got chest x-rays.

Still there are some questions that still need to be answered. For instance, it's not clear whether screening with spiral CT scans would have the same effect on different groups of people, like those who smoked less (or not at all) or younger people. It's also not clear what the best screening schedule might be – how often the scans should be done, how long they should be kept up, *etc.* Also, the lung cancers that were found early were the non-small cell type, so it is not yet clear how to find small cell lung cancer early.

Spiral CT scans are also known to have some downsides that need to be taken into account. One drawback of this test is that it also finds a lot of things that turn out not to be cancer but that still need to be tested to be sure. For some people, this may lead to further, sometimes unnecessary, tests such as CT scans, or even more invasive tests such as biopsies or surgery. Spiral CT scans also expose people to a small amount of radiation with each test. While it is less than the dose from a standard CT, it is more than the dose for a chest x-ray.

These factors, and others, need to be taken into account by people and their doctors who are thinking about whether screening with spiral CT scans is right for them.

Current screening recommendations

Although the American Cancer Society has not yet developed lung cancer screening guidelines, it has plans to do so in the future. In the meantime, some people who are at

higher risk (and their doctors) may want to think about whether screening is right for them.

While a full cancer screening guideline is being developed, the American Cancer Society has some guidance about screening for people who are concerned about their risk of lung cancer. This guidance is complex; it is for patients and their doctors. The screening guidance depends mainly on a person's age and smoking history.

The full text of this guidance can be found in the document, *American Cancer Society Interim Guidance on Lung Cancer Screening*. A brief summary of this guidance, can be found in the "Can small cell lung cancer be found early?" section of our *Lung Cancer detailed guide*. You may also want to talk to your doctor about these recommendations to see if screening might be right for you.

Even with the promising results from the NLST, people who smoke should keep in mind that the best way to avoid dying from lung cancer is to stop smoking. For help quitting smoking, see our *Guide to Quitting Smoking* or call the American Cancer Society at 1-800-227-2345.

Common signs and symptoms of lung cancer

Most lung cancers do not cause symptoms until they have spread, but you should report any of the following problems to your doctor right away. Often these problems are caused by something other than cancer. If lung cancer is found, getting treatment right away might mean your cancer could be cured. Or you could live longer with a better quality of life. The most common symptoms of lung cancer are:

- A cough that does not go away or gets worse
- Chest pain, often made worse by deep breathing, coughing, or laughing
- Hoarseness
- Weight loss and loss of appetite
- Coughing up bloody or rust-colored sputum (spit or phlegm)
- Shortness of breath
- Feeling weak or tired
- Infections such as bronchitis and pneumonia that don't go away or keep coming back
- New onset of wheezing

When lung cancer spreads to distant organs, it may cause:

- Bone pain (like pain in the back or hips)
- Weakness or numbness of the arms or legs
- Headache, dizziness, balance problems, or seizure

- Yellow coloring of the skin and eyes (jaundice)
- Lumps near the surface of the body, caused by cancer spreading to the skin or to lymph nodes in the neck or above the collarbone

If you have any of these symptoms, you should see a doctor right away.

Some lung cancers can cause a group of symptoms called *syndromes*.

Horner syndrome

Cancers of the top part of the lungs (sometimes called *Pancoast tumors*) may damage a nerve that passes from the upper chest into your neck. This can cause severe shoulder pain. Sometimes these tumors also cause a group of symptoms called Horner syndrome:

- Drooping or weakness of one eyelid
- Having a smaller pupil (dark part in the center of the eye) in the same eye
- Reduced or absent sweating on the same side of the face

Conditions other than lung cancer can also cause Horner syndrome.

Superior vena cava syndrome

The superior vena cava (SVC) is a large vein that carries blood from the head and arms back to the heart. It passes next to the upper part of the right lung and the lymph nodes inside the chest. Tumors in this area may push on the SVC, which can cause the blood to back up in the veins. This can cause swelling in the face, neck, arms, and upper chest (sometimes with a bluish-red skin color). It can also cause headaches, dizziness, and a change in mental function if it affects the brain. While SVC syndrome can come on slowly over time, in some cases it can become life-threatening, and needs to be treated right away.

Paraneoplastic syndromes

Some lung cancers may make hormone-like substances that enter the bloodstream and cause problems with other tissues and organs, even though the cancer has not spread to those tissues or organs. These problems are called *paraneoplastic syndromes*. Sometimes these syndromes may be the first symptoms of lung cancer. Because the symptoms affect other organs, patients and their doctors may suspect at first that something other than lung cancer is causing them.

Most of the symptoms listed here are more likely to be caused by something other than lung cancer. Still, if you have any of these problems, you should see a doctor right away.

If your doctor thinks you might have lung cancer

After asking questions about your health and doing a physical exam, your doctor might want to do some of the following tests:

Imaging tests

There are a number of different tests that can make pictures of the inside of your body. Some of these are used to find lung cancer, to see if it has spread, to find out whether treatment is working, or to spot a cancer that has come back after treatment.

Chest x-ray: This is often the first test your doctor will do to look for any spots on the lungs. It is a plain x-ray of your chest. If the x-ray is normal, you most likely do not have lung cancer. If anything does not look normal, the doctor may order more tests.

CT scan (computed tomography): A CT (or CAT) scan uses x-rays to produce detailed cross-sectional pictures of your body. The CT scanner takes many pictures as it moves around you. A computer then combines these pictures into a detailed picture of a slice of your body.

Before the CT scan, you may be asked to drink a special liquid or you may have an IV (intravenous) line through which you are given a "dye." This helps better outline structures in your body. The dye may cause some flushing (a feeling of warmth, especially in the face). Some people are allergic and get hives. Rarely, more serious problems like trouble breathing or low blood pressure can happen. Be sure to tell the doctor if you have ever had a reaction to any contrast material used for x-rays.

CT scans take longer than regular x-rays. You need to lie still on a table while they are being done. You might feel a bit confined by the ring-shaped machine you have to lie in while the pictures are being taken.

The CT scan will give the doctor exact information about the size, shape, and place of a tumor. It can also help find swollen lymph nodes that might contain cancer. CT scans are also used to find tumors in other organs that might be the spread of lung cancer.

In cases where the doctor suspects that the cancer lies deep within the body, a CT scan can be used to guide a biopsy needle right into the that place. For this, you stay on the CT scanning table while the doctor moves a biopsy needle through the skin and toward the mass. Once the doctor can see that the needle is within the mass, a piece of it is removed and looked at under a microscope.

MRI scan (magnetic resonance imaging): Like CT scans, MRI scans give detailed pictures of soft tissues in the body. But MRI scans use radio waves and strong magnets instead of x-rays. MRI scans take longer than x-rays – often up to an hour. Also, you have to be placed inside a tube-like machine, which upsets some people. Newer, open MRI machines can sometimes help with this if needed. MRI scans are useful in finding lung cancer that has spread to the brain or spinal cord.

PET scan (positron emission tomography): For a PET scan, a form of radioactive sugar is injected into the blood. Cancer cells in the body take in large amounts of the sugar. A special camera can then spot the radioactivity. This test can show whether the cancer has spread to the lymph nodes or other parts of the body. It is also helpful in telling whether a spot on your chest x-ray is cancer. Some machines combine a CT and a PET scan to even

better pinpoint tumors. This is the type of PET most often used in small cell lung cancer patients.

Bone scan: For a bone scan a small amount of radioactive substance is put into your vein. The amount used is very low and it causes no long-term effects. This substance builds up in areas of bone that may not be normal because of cancer. These will be seen on the bone scan pictures as dense, gray to black areas, called "hot spots." While these areas may suggest the presence of metastatic cancer, other problems can also cause hot spots.

Bone scans are done mainly when there is reason to think the cancer may have spread to the bones (because of symptoms such as bone pain) and other test results aren't clear. PET scans can usually show the spread of cancer to bones, so bone scans aren't usually needed if a PET scan has already been done.

Other tests to used to find lung cancer and its spread

The tests described below can be used to be sure that something seen on an imaging test is really lung cancer. These tests are also used to find out the exact type of lung cancer and how far it may have spread.

Bronchoscopy: A lighted, flexible tube (called a bronchoscope) is passed through the mouth or nose and into the larger airways of the lungs. The mouth and throat are sprayed first with a numbing medicine. You may also be given medicine through an intravenous (IV) line to make you feel relaxed. This test can help see tumors, or it can be used to take samples of tissue or fluids to see if cancer cells are present.

Endobronchial ultrasound: Ultrasound is a test that uses sound waves to make pictures of parts of your body. For endobronchial ultrasound, a bronchoscope is fitted with an ultrasound device at its tip and is passed down into the windpipe to look at nearby lymph nodes and other structures in the chest. This is done with numbing medicine and light sedation. If areas of concern (such as swollen lymph nodes) are seen on the ultrasound, a hollow needle can be passed through the bronchoscope and guided by ultrasound into the area to take biopsy samples. The samples are then looked at under a microscope to see if there are cancer cells.

This test may be used if the doctor is thinking about surgery as a part of treatment, which is not often the case for small cell lung cancer.

Endoscopic esophageal ultrasound (EUS): This test is much like an endobronchial ultrasound, except an endoscope (a lighted, flexible tube) is used. It is passed down the throat and into the esophagus (the swallowing tube that connects the throat to the stomach). The esophagus lies just behind the windpipe. This test is done with numbing medicine and drugs to help you relax (this is called light sedation).

Ultrasound images taken from inside the esophagus can help find large lymph nodes inside the chest that might contain lung cancer. If areas of concern (such as swollen lymph nodes) are seen on the ultrasound, a hollow needle can be passed through the

endoscope to get biopsy samples of them. The samples are then looked at under a microscope to see if they contain cancer cells.

This test may be used if the doctor is thinking about surgery as a part of treatment, which is not often the case for small cell lung cancer.

Mediastinoscopy and mediastinotomy: Both of these tests may be done so the doctor can look at and take samples of the structures in the area between the lungs (this area is called the *mediastinum*). They are done in an operating room while you are in a deep sleep (under general anesthesia). The main difference between the two is in the place and size of the cut (incision) needed to look into this area.

Thoracentesis: This test is done to check whether fluid around the lungs is caused by cancer or by some other problem, such as heart failure or an infection. First, the skin is numbed and then a hollow needle is placed between the ribs to drain the fluid. The fluid is checked for cancer cells.

Thoracoscopy: For this test, drugs are used to put you to sleep, and a small cut is made in your chest. The doctor then uses a thin, lighted tube connected to a video camera and screen to look at the space between the lungs and the chest wall. By doing this, the doctor can see any cancer deposits on the lung or the lining of the chest wall and take out small pieces of tissue to be looked at under the microscope. Thoracoscopy can also be used to get samples of lymph nodes and fluid and to tell whether a tumor is growing into nearby tissues or organs.

Bone marrow aspiration and biopsy: These two tests are usually done at the same time. For these tests you lie on your side or on your belly. The skin over the back of your hip is cleaned. After the area is numbed, a thin, hollow needle is inserted into the hip bone to suck out (aspirate) a small amount of fluid from the marrow. A larger needle is then used to remove a small piece of the hip bone and some marrow. Even with the numbing medicine, most patients still have some brief pain when the samples are removed. The samples are then checked for cancer cells. This is done in some patients to be sure that the cancer has not spread to the bone marrow.

Sampling tissues and cells

Symptoms and the results of imaging tests may strongly suggest that lung cancer is present, but the final diagnosis of lung cancer is made by looking at lung cells under a microscope. The cells can be obtained in different ways. One or more of the tests below may be used to find out if a lung mass seen on imaging tests is indeed lung cancer. These tests can also be used to tell the exact type of lung cancer you have and to help figure out how far it may have spread.

A doctor (called a pathologist) who is an expert in using lab tests to diagnose diseases like cancer will look at the cells under a microscope. The results will be described in a pathology report, which should take about a week. If you have any questions about your results or any other tests, be sure and ask your doctor. If needed, you can get a second opinion by having your tissue sample sent to a pathologist at another lab.

Sputum cytology: A sample of sputum (mucus you cough up from the lungs) is looked at under a microscope to see if cancer cells are present.

Fine needle biopsy (FNA): For this test, a long, thin (fine) needle is put into the place in the lung that might be cancer to remove a sample of cells. (The skin where the needle is to be put in may first be numbed with medicine.) The sample is looked at in the lab to see whether there are cancer cells. An FNA biopsy may also be done to take samples of lymph nodes around the windpipe (trachea) and the larger tubes that carry air to the lungs (bronchi). (In some cases, if the results aren't clear, a larger needle may be used to remove a slightly bigger piece of lung tissue. This is known as a *core needle biopsy*.)

Sometimes, air may leak out of the lung at the biopsy site and into the space between the lung and the chest wall. This can cause part of the lung to collapse and may cause trouble with breathing. This often gets better without any treatment. If not, a small tube is put into the chest space and the air is sucked out over a day or two, after which it should heal on its own.

Lab tests of biopsy and other samples

Samples from biopsies or other tests are sent to a lab. There, a doctor looks at the samples under a microscope to find out if they contain cancer and if so, what type of cancer it is. Special tests may be needed to help classify the cancer. Cancers from other organs can spread to the lungs, so it is very important to find out where the cancer started so the right treatment can be given.

Blood tests

Blood tests are not used to find lung cancer, but they are done to get a sense of a person's overall health. A complete blood count (CBC) shows whether your blood has normal numbers of different cell types. This test will be done often if you are treated with chemotherapy because these drugs can affect the blood-forming cells of the bone marrow. Other blood tests can spot problems in different organs like the kidneys, liver, and bones.

Pulmonary function tests

There are different types of pulmonary function tests but they all basically involve having you breathe in and out through a tube that is connected to different machines. As a rule, they are only needed if surgery might be an option in treating the cancer. Since surgery is rarely used to treat small cell lung cancer, these tests are not often done for patients known to have small cell lung cancer.

Staging for small cell lung cancer

Staging is the process of finding out how far the cancer has spread. This is very important because your treatment and your outlook (prognosis) depend on the stage of the cancer.

The tests described in the section, "How is small cell lung cancer found?" are also used to stage lung cancer.

A staging system is a standard way for doctors to describe how large a cancer is and how far it has spread. There are 2 types of staging.

The *clinical stage* is based on the results of the physical exam, biopsies, and imaging tests (CT scan, chest x-ray, PET scan, *etc.*).

If you have surgery, your doctor can also determine a *pathologic stage*, which is based on the same factors as the clinical stage, plus what is found during the surgery.

The clinical and pathologic stages may be different in some cases. During surgery the doctor may find cancer in a place that did not show up on imaging tests. That could mean a more advanced pathologic stage. Because most patients with lung cancer do not have surgery, the clinical stage is most often used when describing the cancer.

Small cell lung cancer staging systems

There are 2 staging systems that can be used to describe the spread of small cell lung cancer.

Limited vs. extensive stage

For small cell lung cancer a 2-stage system is most often used. This system divides small cell lung cancers into *limited stage* or *extensive stage*.

Limited stage usually means that the cancer is only in one lung and perhaps in lymph nodes on the same side of the chest. The cancer is most often confined to an area that is small enough to be treated with radiation.

If the cancer has spread to the other lung, to lymph nodes on the other side of the chest, or to distant organs, it is called extensive. Many doctors also call cancer that has spread to the fluid around the lung extensive stage.

Small cell lung cancer is staged in this way because it helps to separate people who may be helped by treatments such as surgery and radiation therapy to try to cure the cancer (limited stage) from those for whom these treatments aren't likely to cure the cancer (extensive stage).

The TNM staging system

A more formal system to describe the growth and spread of lung cancer is the American Joint Committee on Cancer (AJCC) TNM staging system. Stages are described using Roman numerals from 0 to IV (0 to 4). Some stages are further divided into A and B. As a rule, the lower the number, the less the cancer has spread. A higher number, such as stage IV (4), means a more advanced cancer.

This system is used more often for non-small cell lung cancer. It is used less often for small cell lung cancer, mainly because treatment options don't vary much between these detailed stages.

Survival rates for small cell lung cancer

Some people with cancer may want to know the survival rates for their type of cancer. Others may not find the numbers helpful, or may even not want to know them. If you do not want to know them, stop reading here and skip to the next section.

Survival rates are a way for doctors and patients to get a general idea of the outlook for people with a certain type and stage of cancer. The 5-year survival rate refers to the percentage of patients who live at least 5 years after their cancer is found. Of course, many people live much longer than 5 years. Five-year *relative* survival rates compare the number of people who are still alive 5 years after their cancer was found to the survival of others the same age who don't have cancer. This is a better way to see the impact that cancer can have on survival.

The numbers below are 5-year relative survival rates from the National Cancer Institute's Surveillance, Epidemiology and End Results (SEER) database. They are based on people with small cell lung cancer treated between 1988 and 2001. Improvements in treatment since then mean that the survival rates for people with these cancers may now be higher.

Stage	5-year relative survival rate
I	31%
II	19%
III	8%
IV	2%

While these numbers provide an overall picture, keep in mind that every person's situation is unique and the statistics can't predict exactly what will happen in your case. Talk with your cancer care team if you have questions about your own chances of a cure, or how long you might survive your cancer. They know your situation best.

How is small cell lung cancer treated?

This information represents the views of the doctors and nurses serving on the American Cancer Society's Cancer Information Database Editorial Board. These views are based on their interpretation of studies published in medical journals, as well as their own professional experience.

The treatment information in this document is not official policy of the Society and is not intended as medical advice to replace the expertise and judgment of your cancer care team. It is intended to help you and your family make informed decisions, together with your doctor.

Your doctor may have reasons for suggesting a treatment plan different from these general treatment options. Don't hesitate to ask him or her questions about your treatment options.

Choosing a treatment plan for small cell lung cancer

If you have small cell lung cancer, the main treatment will most likely be chemotherapy (chemo), either alone or with radiation. Very rarely, surgery might be done if it is limited stage cancer.

After the cancer is found and staged, your doctor will talk to you about treatment choices. Give yourself time to take in what you have learned. The most important things to think about include the stage and type of cancer, your overall health, the likely side effects of the treatment, and the chance of curing the cancer or helping you live longer. Age alone should not keep you from having treatment. Older people can be helped by treatment as much as younger people as long as their general health is good.

You may have different types of doctors on your treatment team, depending on the stage of your cancer and your treatment options. Many other experts may be involved in your care as well (nurses, respiratory therapists, social workers, *etc.*)

Surgery for small cell lung cancer

Surgery is rarely used as the main treatment in small cell lung cancer. In fewer than 1 out of 20 cases, the cancer is found as only one tumor, with no spread to lymph nodes or other organs. In these cases, surgery might be helpful. If used, it is usually followed by treatment with chemo and maybe radiation.

There are different operations that can be used to treat small cell lung cancer. For each of them you will be in a deep sleep (under general anesthesia) and will have a cut (surgical incision) between the ribs in the side of the chest.

- Pneumonectomy: the entire lung is removed.
- Lobectomy: a section (lobe) of the lung is removed.
- Segmentectomy or wedge resection: part of a lobe is removed.
- Sleeve resection: a section of a large airway is removed and the lung is reattached.

As a rule, lobectomy is the preferred operation for small cell lung cancers treated with surgery. With any of these operations, lymph nodes are also removed to look for possible spread of the cancer.

You will most likely need to spend about a week in the hospital after the surgery. There will be some pain because the surgeon has to cut through the ribs to get to the lungs. Other possible problems include bleeding, infections, and pneumonia.

Video-assisted thoracic surgery (VATS): This is a new kind of surgery for some people with early stage lung cancer. A tiny camera can be put through a small hole in the chest to help the surgeon see the tumor. One or 2 other small cuts (incisions) are made in the skin, and long instruments are passed through these to remove the tumor. Since only small cuts are needed, there is a shorter hospital stay and less pain after surgery. This approach is most often used for tumors smaller than about 1½ inches near the outside of the lung. The cure rate seems to be the same as for standard surgery. The doctor who does this surgery should have experience because it takes a great deal of skill.

Possible side effects of surgery

Problems after surgery depend on how much surgery was done and a person's health beforehand. Serious problems can include a lot of bleeding, infection, and pneumonia. While it is rare, in some cases people may not survive the surgery, which is why it is very important that surgeons select patients with care.

People whose lungs are in good shape (other than the cancer) can often return to normal activities after some time if a lobe or even an entire lung is removed. But if they also have problems like emphysema or chronic bronchitis, which are common in heavy smokers, they may have long-term shortness of breath.

The surgeon must spread ribs to get to the lung when doing a thoracotomy, so the incision will hurt for some time after surgery. Your activity will be limited for at least a month.

Surgery and other methods to relieve problems

In some cases, surgery or other treatments may be used to help with the symptoms of the cancer (rather than trying to remove all of the cancer). For instance, laser surgery can be used to open an airway blockage that may be causing pneumonia or shortness of breath. Sometimes, metal or plastic tubes called stents may be placed in the airways to help keep them open. Other methods like radiation may also be used.

Sometimes fluid collects in the chest and makes it hard to breathe. This fluid can be removed through a small tube placed in the chest. After the fluid is drained out, either talc or some type of drug is placed into the chest. This will start a reaction that will help seal the space and prevent future fluid build-up.

Radiation treatment for small cell lung cancer

Radiation therapy is treatment with high-energy rays (like x-rays) to kill cancer cells or shrink tumors. The radiation comes from outside the body (external radiation). In small cell lung cancer it is most often used along with chemotherapy (in limited stage disease) to treat the tumor and lymph nodes in the chest.

Radiation might be used on the brain to try to prevent the spread of cancer there. Radiation can also be used to relieve symptoms such as pain, bleeding, trouble swallowing, or problems caused by the cancer spreading to the brain.

Each treatment lasts only a few minutes, although the setup time – getting you into place for treatment – takes longer. Most often, radiation treatments are given once or twice a day, 5 days a week for several weeks. Radiation to relieve symptoms or to prevent spread to the brain is given for shorter periods of time. Standard radiation is used less often than in the past. Newer methods such as 3D-CRT and IMRT allow doctors to be more precise in treating lung cancers while reducing radiation to nearby healthy tissues.

Possible side effects of radiation treatment

- Sunburn-like skin problems
- Hair loss (in the place where the radiation enters the body)
- Tiredness (fatigue)
- Nausea and vomiting
- Loss of appetite and weight loss

Radiation can affect the blood-forming cells in the bone marrow. This can lead to low blood counts. This can lead to:

- Increased chance of infections (from low white blood cell counts)
- Easy bruising or bleeding (from low blood platelet counts)
- Fatigue (from low red blood cell counts)

If your esophagus, which is in the middle of your chest, is exposed to radiation, it could cause a sore throat and trouble swallowing during treatment. This may make it hard to eat anything other than soft foods or liquids for a while.

When chemotherapy is given with radiation, many of the side effects are worse. Most side effects improve or even go away after treatment ends.

Side effects of radiation therapy to the brain usually become most serious 1 or 2 years after treatment. These side effects could include memory loss, headaches, trouble with thinking, and less sexual desire. These side effects, though, are usually minor compared to those caused by lung cancer tumors that have spread to the brain.

For more information about radiation treatment, please see our document, *Understanding Radiation Therapy: A Guide for Patients and Families*.

Chemotherapy for small cell lung cancer

Chemotherapy (chemo) is treatment with drugs given into a vein or taken by mouth. These drugs enter the bloodstream and go throughout the body. This treatment is useful for cancer that has spread (metastasized) to organs beyond the lung. Chemo is usually the main treatment for small cell lung cancer.

Doctors give chemo in cycles, with each round of treatment followed by a rest period to allow the body time to recover. Chemo cycles often last about 3 to 4 weeks, and the first round of treatments is most often 4 to 6 cycles. Chemo is not often used for patients in poor health, but older age by itself doesn't mean you can't get chemo.

Possible side effects

Chemo drugs kill cancer cells but they also damage some normal cells, causing side effects. These side effects depend on the type of drugs used, the amount given, and the length of treatment. You could have some of these short-term side effects:

- Hair loss
- Mouth sores
- Loss of appetite
- Nausea and vomiting
- Diarrhea or constipation
- Increased chance of infections (from low white blood cell counts)
- Easy bruising or bleeding (from low blood platelet counts)
- Feeling very tired all the time, called fatigue (from low red blood cell counts)

Some chemo drugs can damage nerves. This can cause numbness in the fingers and toes, and sometimes the arms and legs may feel weak. For find out more, see our document, *Peripheral Neuropathy Caused by Chemotherapy*.

Most side effects go away when treatment is over, but some can last a long time. Be sure to tell your doctor or nurse if you have any side effects, as there are often ways to help. In some cases, the doses of the chemo drugs may need to be lowered or treatment may need to be delayed or stopped to prevent the side effects from getting worse.

To learn more about chemo please see our document, *Understanding Chemotherapy: A Guide for Patients and Families*.

Clinical trials for small cell lung cancer

You may have had to make a lot of decisions since you've been told you have cancer. One of the most important decisions you will make is deciding which treatment is best for you. You may have heard about clinical trials being done for your type of cancer. Or maybe someone on your health care team has mentioned a clinical trial to you.

Clinical trials are carefully controlled research studies that are done with patients who volunteer for them. They are done to get a closer look at promising new treatments or procedures.

If you would like to take part in a clinical trial, you should start by asking your doctor if your clinic or hospital conducts clinical trials. You can also call our clinical trials matching service for a list of clinical trials that meet your medical needs. You can reach this service at 1-800-303-5691 or on our Web site at www.cancer.org/clinicaltrials. You can also get a list of current clinical trials by calling the National Cancer Institute's Cancer Information Service toll-free at 1-800-4-CANCER (1-800-422-6237) or by visiting the NCI clinical trials Web site at www.cancer.gov/clinicaltrials.

There are requirements you must meet to take part in any clinical trial. If you do qualify for a clinical trial, it is up to you whether or not to enter (enroll in) it.

Clinical trials are one way to get state-of-the-art cancer treatment. They are the only way for doctors to learn better methods to treat cancer. Still, they are not right for everyone.

You can get a lot more information on clinical trials in our document called *Clinical Trials: What You Need to Know*. You can read it on our Web site or call our toll-free number and have it sent to you.

Complementary and alternative therapies for small cell lung cancer

When you have cancer you are likely to hear about ways to treat your cancer or relieve symptoms that your doctor hasn't mentioned. Everyone from friends and family to Internet groups and Web sites may offer ideas for what might help you. These methods can include vitamins, herbs, and special diets, or other methods such as acupuncture or massage, to name a few.

What are complementary and alternative therapies?

It can be confusing because not everyone uses these terms the same way, and they are used to refer to many different methods. We use *complementary* to refer to treatments that are used *along with* your regular medical care. *Alternative* treatments are used *instead of* a doctor's medical treatment.

Complementary methods: Most complementary treatment methods are not offered as cures for cancer. Mainly, they are used to help you feel better. Some examples of methods that are used along with regular treatment are meditation to reduce stress, acupuncture to help relieve pain, or peppermint tea to relieve nausea. Some complementary methods are known to help, while others have not been tested. Some have been proven not to be helpful, and a few are even harmful.

Alternative treatments: Alternative treatments may be offered as cancer cures. These treatments have not been proven safe and effective in clinical trials. Some of these methods may be harmful, or have life-threatening side effects. But the biggest danger in most cases is that you may lose the chance to be helped by standard medical treatment. Delays or interruptions in your medical treatments may give the cancer more time to grow and make it less likely that treatment will help.

Finding out more

It is easy to see why people with cancer think about alternative methods. You want to do all you can to fight the cancer, and the idea of a treatment with few or no side effects sounds great. Sometimes medical treatments like chemotherapy can be hard to take, or they may no longer be working. But the truth is that most of these alternative methods have not been tested and proven to work in treating cancer.

As you think about your options, here are 3 important steps you can take:

- Look for "red flags" that suggest fraud. Does the method promise to cure all or most cancers? Are you told not to have regular medical treatments? Is the treatment a "secret" that requires you to visit certain providers or travel to another country?
- Talk to your doctor or nurse about any method you are thinking of using.
- Contact us at 1-800-227-2345 to learn more about complementary and alternative methods in general and to find out about the specific methods you are looking at.

The choice is yours

Decisions about how to treat or manage your cancer are always yours to make. If you want to use a non-standard treatment, learn all you can about the method and talk to your doctor about it. With good information and the support of your health care team, you may be able to safely use the methods that can help you while avoiding those that could be harmful.

What are some questions I can ask my doctor about small cell lung cancer?

As you cope with cancer and cancer treatment, we encourage you to have honest, open talks with your doctor. Feel free to ask any question that's on your mind, no matter how small it might seem. Here are some questions you might want to ask. Take them with you to your next doctor visit. Be sure to add your own questions as you think of them. Nurses, social workers, and other members of the treatment team may also be able to answer many of your questions.

- Would you please write down the exact type of lung cancer I have?
- May I have a copy of my pathology report?
- Has my cancer spread beyond the place where it started?
- What is the stage of my cancer? What does that mean in my case?
- Are there other tests that need to be done before we can decide on treatment?
- Are there other doctors I need to see?

- How much experience do you have treating this type of cancer?
- What treatment choices do I have?
- What do you suggest and why?
- What is the goal of this treatment?
- How long will treatment last? What will it involve? Where will it be done?
- What are my chances of long-term survival, based on my cancer as you see it?
- What risks or side effects are there to the treatment you suggest? How long are they likely to last?
- How quickly do we need to decide on treatment?
- What are the chances of the cancer coming back after treatment? What would we do if that happens?
- Will I lose my hair? If so, what can I do about it?
- What should I do to get ready for treatment?
- What will we do if the treatment doesn't work or if the cancer comes back?
- What type of follow-up will I need after treatment?

Add your own questions below:

Moving on after treatment for small cell lung cancer

For some people with lung cancer, treatment may remove or destroy the cancer. While it can feel good to be done with treatment, it can also be stressful. You may find that you now worry about the cancer coming back. This is a very common concern among those who have had cancer. (When cancer comes back, it is called a *recurrence*.)

It may take a while before your recovery begins to feel real and your fears are somewhat relieved. You can learn more about what to look for and how to learn to live with the chance of cancer coming back in *Living With Uncertainty: The Fear of Cancer Recurrence*.

But for other people, the lung cancer may never go away completely. You may keep on getting treatments with chemo, radiation, or other treatments to help keep the cancer in check. Learning to live with cancer as a kind of chronic disease can be hard and very stressful. Our document, *When Cancer Doesn't Go Away*, talks more about this.

Follow-up care

During and after treatment, your doctors will still want to watch you closely. It is very important to keep all follow-up appointments. During these visits, your doctors will ask about symptoms, do physical exams, and may order blood tests, CT scans, or x-rays.

In people who show no signs of cancer, most doctors recommend follow-up visits about every 2 to 3 months for the first year after treatment, every 3 to 6 months for the next several years, then at least yearly after 5 years. Follow-up is needed to check for cancer that has come back or spread, as well as possible side effects of certain treatments. This is the time for you to ask your health care team any questions you might have and discuss any of your concerns.

Almost any cancer treatment can have side effects. Some may last for a few weeks or months, but others can last the rest of your life. Please tell your cancer care team about any symptoms or side effects that bother you so they can help you manage them.

It is important to keep health insurance. Tests and doctor visits cost a lot, and even though no one wants to think of their cancer coming back, this could happen.

Should your cancer come back, our document *When Your Cancer Comes Back: Cancer Recurrence* helps you manage and cope with this phase of your treatment.

Keep your health insurance and copies of your medical records

At some point after your cancer is found and treated, you may find yourself in the office of a new doctor. It is important that you be able to give your new doctor the exact details of your diagnosis and treatment. Make sure you have this information handy and always keep copies for yourself:

- A copy of your pathology report from any biopsy or surgery
- If you had surgery, a copy of your operative report
- If you were in the hospital, a copy of the discharge summary that the doctor wrote when you were sent home
- If you had radiation treatment, a copy of the treatment summary
- If you had chemo, a list of your drugs, drug doses, and when you took them
- Copies of your x-rays, CT scans, and other imaging tests (these can often be put on a DVD)

The doctor may want copies of this information for his records, but always keep copies for yourself.

Lifestyle changes after treatment for small cell lung cancer

Having cancer and dealing with treatment can take a lot of time and energy, but it can also be a time to look at your life in new ways. Maybe you are thinking about how to improve your health over the long term.

Make healthier choices

For many people, a diagnosis of cancer helps them focus on their health in ways they may not have thought much about in the past. Are there things you could do that might make you healthier? Maybe you could try to eat better or get more exercise. Maybe you could cut down on the alcohol, or give up tobacco. Even things like keeping your stress level under control may help. Now is a good time to think about making changes that can have positive effects for the rest of your life. You will feel better and you will also be healthier.

You can start by working on those things that worry you most. Get help with those that are harder for you. For instance, if you smoke, one of the best things you can do to improve your chances for treatment success is to quit. Studies have shown that patients who stop smoking after a finding of lung cancer have better outcomes than those who don't. Quitting can help improve lung function and have a host of other health benefits as well. If you are thinking about quitting smoking and need help, call us at 1-800-227-2345.

Eating better

Eating right is hard for many people, but it can be even harder to do during and after cancer treatment. Treatment may change your sense of taste. Nausea can be a problem. You may not feel like eating and lose weight when you don't want to. Or you may have gained weight that you can't seem to lose. All of these things can be very frustrating.

If treatment caused weight changes or eating or taste problems, do the best you can and keep in mind that these problems usually get better over time. You may find it helps to eat small portions every 2 to 3 hours until you feel better. You may also want to ask your cancer team about seeing a dietitian, an expert in nutrition who can give you ideas on how to deal with these treatment side effects.

One of the best things you can do after cancer treatment is put healthy eating habits into place. You may be surprised at the long-term benefits of some simple changes, like increasing the variety of healthy foods you eat. Getting to and staying at a healthy weight, eating a healthy diet, and limiting your alcohol intake may lower your risk for some other types of cancer, as well as having many other health benefits.

Rest, fatigue, and exercise

Feeling tired (fatigue) is a very common problem during and after cancer treatment. This is not a normal type of tiredness but a "bone-weary" exhaustion that doesn't get better with rest. For some people, fatigue lasts a long time after treatment and can keep them

from staying active. But exercise can actually help reduce fatigue and the sense of depression that sometimes comes with feeling so tired.

If you are very tired, though, you will need to balance activity with rest. It is OK to rest when you need to. To learn more about fatigue, please see our documents, *Fatigue in People With Cancer* and *Anemia in People With Cancer*.

If you were very ill or weren't able to do much during treatment, it is normal that your fitness, staying power, and muscle strength declined. You need to find an exercise plan that fits your own needs. Talk with your health care team before starting. Get their input on your exercise plans. Then try to get an exercise buddy so that you're not doing it alone.

Exercise can improve your physical and emotional health.

- It improves your cardiovascular (heart and circulation) fitness.
- It makes your muscles stronger.
- It reduces fatigue.
- It can help lower anxiety and depression.
- It can make you feel generally happier.
- It helps you feel better about yourself.

And long term, we know that getting regular physical activity plays a role in helping to lower the risk of some cancers, as well as having other health benefits.

Can I lower my risk of the cancer growing or coming back?

Most people want to know whether there are lifestyle changes they can make to reduce their risk of the cancer growing or coming back. Unfortunately, for most cancers there isn't much solid evidence to guide people. This doesn't mean that nothing will help – it's just that for the most part this isn't something that has been well studied. Most studies have looked at lifestyle changes as ways to prevent cancer in the first place, not slowing it down or keeping it from coming back.

But there are some things people can do that might help them live longer or reduce the risk of lung cancer coming back.

Quitting smoking: If you smoke, quitting is important. It has been shown to help improve outcomes and reduce the risk of the cancer coming back, especially in people with early stage lung cancer. Of course, quitting smoking may have other health benefits as well, like lowering the risk of some other cancers. If you need help quitting, talk to your doctor or call the American Cancer Society at 1-800-227-2345.

What you eat: Possible links between diet and lung cancer growing or coming back are much less clear. Because of the lack of data in this area, it's important to talk with your health care team before making any major changes to your (including taking any supplements) to try to improve your outlook.

How about your emotional health after small cell lung cancer?

Once your treatment ends, you may be surprised by the flood of emotions you go through. This happens to a lot of people. You may find that you think about the effect of your cancer on things like your family, friends, and career. Money may be a concern as the medical bills pile up. Unexpected issues may also cause concern – for instance, as you get better and need fewer doctor visits, you will see your health care team less often. This can be hard for some people.

This is a good time to look for emotional and social support. You need people you can turn to. Support can come in many forms: family, friends, cancer support groups, church or spiritual groups, online support communities, or private counselors.

The cancer journey can feel very lonely. You don't need to go it alone. Your friends and family may feel shut out if you decide not to include them. Let them in – and let in anyone else who you feel may help. If you aren't sure who can help, call your American Cancer Society at 1-800-227-2345 and we can put you in touch with a group or resource that may work for you.

You can't change the fact that you have had cancer. What you can change is how you live the rest of your life -- making healthy choices and helping your body and mind feel well.

If treatment for small cell lung cancer stops working

When a person has had many different treatments and the cancer has not been cured, over time the cancer tends to resist all treatment. At this time you may have to weigh the possible benefits of a new treatment against the downsides, like treatment side effects and clinic visits.

This is likely to be the hardest time in your battle with cancer – when you have tried everything within reason and it's just not working anymore. Your doctor may offer you new treatment, but you will need to talk about whether the treatment is likely to improve your health or change your outlook for survival.

If you want to keep on getting treatment for as long as you can, you need to think about the odds of treatment having any benefit and how this compares to the possible risks and side effects. In many cases, your doctor can tell you how likely it is the cancer will respond to treatment you are thinking about. For instance, the doctor may say that more treatment might have about a 1 in 100 chance of working. Some people are still tempted to try this. But it is important to think about and understand your reasons for choosing this plan.

No matter what you decide to do, it is important for you to feel as good as possible. Make sure you are asking for and getting treatment for pain, nausea, or any other problems you may have. This type of treatment is called *palliative* treatment. It helps relieve symptoms but is not meant to cure the cancer.

At some point you may want to think about hospice care. Most of the time it is given at home. Your cancer may be causing symptoms or problems that need to be treated. Hospice focuses on your comfort. You should know that having hospice care doesn't mean you can't have treatment for the problems caused by your cancer or other health issues. It just means that the purpose of your care is to help you live life as fully as possible and to feel as well as you can.

You can learn more about this in our documents, *Hospice Care*.

What's new in small cell lung cancer research?

Lung cancer research is going on now in many medical centers throughout the world.

Prevention

Tobacco

At this time, many researchers believe that prevention offers the greatest promise for fighting lung cancer. Smoking still accounts for almost 9 out of 10 lung cancer deaths. This percentage is likely even higher for small cell lung cancers. Studies are going on to look at how best to help people quit smoking through counseling, nicotine replacement, and other medicines. Other studies are looking at ways to convince young people not to start smoking. Researchers are also looking at differences in genes that may make some people much more likely to get lung cancer if they smoke or are exposed to someone else's smoke.

Diet, nutrition, and medicines

Although researchers are looking for ways to use vitamins or medicines to prevent lung cancer in people at high risk, so far, these have not proved to help. Many researchers think that simply following the American Cancer Society nutrition recommendations (such as staying at a healthy weight and eating at least 5 servings of fruits and vegetables each day) may be the best approach.

Finding lung cancer

As mentioned in the section, "How is small cell lung cancer found?" a large study called the National Lung Screening Trial (NLST) recently found that spiral CT scanning in people at high risk of lung cancer (due to smoking history) lowered the risk of death from lung cancer when compared to chest x-rays. Doctors will learn more about what this study means in the near future.

Another approach uses new ways to try to find cancer cells in sputum (spit) samples. Researchers have also found some changes that often affect the DNA of lung cancer

cells. New tests might be able to spot these changes and find lung cancer at an earlier stage.

Fluorescence bronchoscopy (also known as *autofluorescence bronchoscopy*) is a method that may help doctors find some lung cancers earlier, when they could be easier to treat. For this test, the doctor puts a bronchoscope through the mouth or nose and into the lungs. The end of the bronchoscope has a special light on it. The light causes abnormal areas in the airways to show up in a different color than healthy parts of the airway. Some cancer centers now use this to look for early lung cancers, especially when no tumors are seen with normal bronchoscopy.

An imaging test called virtual bronchoscopy uses CT scans to create detailed 3-D pictures of the airways in the lung. The images can be looked at as if the doctor were actually using a bronchoscope. There are benefits and drawbacks to this approach. But it may be a useful tool in some cases, such as in people who might be too sick to get a standard bronchoscopy. This test will likely be used more often as the technology improves.

Treatment

Chemotherapy

Many clinical trials are being done to see how well newer combinations of chemo drugs work. These studies are also looking at ways to reduce side effects, especially in patients who are older and have other health problems. Doctors are also searching for better ways to combine chemo with radiation and other treatments.

Some new chemo drugs have shown promise in early studies and are now being tested in larger clinical trials.

Targeted therapies

Researchers are learning more about the inner workings of lung cancer cells that control their growth and spread. This is being used to develop new targeted therapies. These drugs work in a different way from standard chemo drugs. They often have different (and less severe) side effects. Many of these are being tested in clinical trials to see if they can help people with advanced cancer to live longer or to relieve their symptoms.

Drugs to keep new blood vessels from growing (anti-angiogenesis drugs)

For cancers to grow, new blood vessels must develop to nourish the cancer cells within tumors. This process is called angiogenesis. New drugs that slow or stop angiogenesis are being studied as lung cancer treatments. Some have already been successfully used for other cancer types. For instance, a drug called bevacizumab (Avastin) has been shown to help patients with some types of non-small cell lung cancer, and is now being tested in small cell lung cancer. In a study of small cell lung cancer, it helped stop some of the cancers from growing for a time, but didn't seem to help the patients live longer.

Vaccines

Vaccines that boost the body's immune system to better kill lung cancer cells are being tested in clinical trials. Unlike vaccines against infections like measles or mumps, these vaccines are designed to help treat, not prevent, lung cancer. One possible advantage of these types of treatments is that they seem to have fewer side effects, so they might be useful in people who can't have other treatments. At this time, vaccines are only being used in clinical trials.

More information about small cell lung cancer

From your American Cancer Society

The following information may also be helpful to you. These materials may be ordered from our toll-free number.

Small Cell Lung Cancer Detailed Guide (also in Spanish)

After Diagnosis: A Guide for Patients and Families (also in Spanish)

Caring for the Patient With Cancer at Home: A Guide for Patients and Families (also in Spanish)

Guide to Quitting Smoking (also in Spanish)

Lasers in Cancer Treatment

Living With Uncertainty: The Fear of Cancer Recurrence

Pain Control: A Guide for Patients and Families (also in Spanish)

Peripheral Neuropathy Caused by Chemotherapy

Questions About Smoking, Tobacco, and Health (also in Spanish)

Surgery (also in Spanish)

Understanding Chemotherapy: A Guide for Patients and Families (also in Spanish)

Understanding Radiation Therapy: A Guide for Patients and Families (also in Spanish)

When Your Cancer Comes Back: Cancer Recurrence

The following books are available from the American Cancer Society. Call us at 1-800-227-2345 to ask about costs or to place your order.

American Cancer Society Complete Guide to Complementary & Alternative Cancer Therapies

American Cancer Society Complete Guide to Nutrition for Cancer Survivors

American Cancer Society's Guide to Pain Control

Cancer in the Family: Helping Children Cope With a Parents Illness

Caregiving: A Step-By-Step Resource for Caring for the Person With Cancer at Home

What Helped Get Me Through: Cancer Patients Share Wisdom and Hope

What to Eat During Cancer Treatment

When the Focus Is on Care: Palliative Care and Cancer

National organizations and Web sites*

Along with the American Cancer Society, other sources of information and support include:

American Lung Association

Toll-free number: 1-800-586-4872 (1-800-LUNGUSA)

Web site: www.lungusa.org

Lungcancer.org

Toll-free number: 1-800-813-4673 (1-800-813-HOPE)

Web site: www.lungcancer.org

Lung Cancer Alliance

Toll-free number: 1-800-298-2436 (United States only)

Web site: www.lungcanceralliance.org

National Cancer Institute

Toll-free number: 1-800-422-6237 (1-800-4-CANCER)

Web site: www.cancer.gov

** Inclusion on this list does not imply endorsement by the American Cancer Society.*

No matter who you are, we can help. Contact us anytime, day or night, for information and support. Call us at **1-800-227-2345** or visit www.cancer.org.

Last Medical Review: 3/14/2012

Last Revised: 3/14/2012

2012 Copyright American Cancer Society

For additional assistance please contact your American Cancer Society
1 · 800 · ACS-2345 or www.cancer.org