

Malignant Mesothelioma

What is cancer?

The body is made up of trillions of living cells. Normal body cells grow, divide into new cells, and die in an orderly way. During the early years of a person's life, normal cells divide faster so the person can grow. After the person becomes an adult, most cells divide only to replace worn-out or dying cells or to repair injuries.

Cancer begins when cells in a part of the body start to grow out of control. There are many kinds of cancer, but they all start because of out-of-control growth of abnormal cells.

Cancer cell growth is different from normal cell growth. Instead of dying, cancer cells continue to grow and form new, abnormal cells. Cancer cells can also invade (grow into) other tissues, something normal cells cannot do. Growing out of control and invading other tissues is what makes a cell a cancer cell.

Cells become cancer cells because of damage to DNA. DNA is in every cell and directs all its actions. In a normal cell, when DNA gets damaged the cell either repairs the damage or the cell dies. In cancer cells, the damaged DNA is not repaired, but the cell doesn't die like it should. Instead, this cell goes on making new cells that the body does not need. These new cells will all have the same damaged DNA as the first cell does.

People can inherit damaged DNA, but most DNA damage is caused by mistakes that happen while the normal cell is reproducing or by something in our environment. Sometimes the cause of the DNA damage is something obvious, like cigarette smoking. But often no clear cause is found.

In most cases the cancer cells form a tumor. Some cancers, like leukemia, rarely form tumors. Instead, these cancer cells involve the blood and blood-forming organs and circulate through other tissues where they grow.

Cancer cells often travel to other parts of the body, where they begin to grow and form new tumors that replace normal tissue. This process is called *metastasis*. It happens when the cancer cells get into the bloodstream or lymph vessels of our body.

No matter where a cancer spreads, it is always named for the place where it started. For example, breast cancer that has spread to the liver is still called breast cancer, not liver cancer. Likewise, prostate cancer that has spread to the bone is metastatic prostate cancer, not bone cancer.

Different types of cancer can behave very differently. For example, lung cancer and breast cancer are very different diseases. They grow at different rates and respond to different treatments. That is why people with cancer need treatment that is aimed at their particular kind of cancer.

Not all tumors are cancerous. Tumors that aren't cancer are called benign. Benign tumors can cause problems -- they can grow very large and press on healthy organs and tissues. But they cannot grow into (invade) other tissues. Because they can't invade, they also can't spread to other parts of the body (metastasize). These tumors are almost never life threatening.

What is malignant mesothelioma?

Malignant mesothelioma is a cancer that starts in cells in the linings of certain parts of the body, especially the chest cavity or abdominal cavity..

A layer of specialized cells called *mesothelial cells* lines the inside of the chest, the abdomen, and the space around your heart. These cells also cover the outer surface of most of your internal organs. The lining formed by these cells is called the *mesothelium*.

The mesothelium helps protect your organs by making a special lubricating fluid that allows organs to move. For example, this fluid makes it easier for your lungs to move (expand and contract) inside the chest when you breathe. The mesothelium has different names in different parts of the body:

- The *pleura* coats the lungs and the cavity containing the lungs in the chest.
- The *peritoneum* coats the abdominal cavity and many of the organs within that cavity.
- The *tunica vaginalis* coats the testicles.
- The *pericardium* coats the heart and creates the cavity that holds the heart in the chest.

Mesothelial tumors can be non-cancerous (benign) or cancerous (malignant).

Malignant mesothelioma

A cancerous tumor of the mesothelium is called a *malignant mesotheliuma*, although this is often shortened to just mesotheliuma. Mesotheliumas can start in 4 main areas in the body.

• **Pleural mesotheliomas** start in the chest. About 3 out of 4 mesotheliomas are pleural mesotheliomas.

- **Peritoneal mesotheliomas** begin in the abdomen. They make up most of the remaining cases.
- Pericardial mesotheliomas start in the covering around the heart and are very rare.
- Mesotheliomas of the tunica vaginalis are very rare tumors that start in the covering layer of the testicles.

Malignant mesotheliomas can also be classified into 3 main types based on how the cells are arranged:

- About 50% to 60% of mesotheliomas are **epithelioid**. This type tends to have a better outlook (prognosis) than the other types.
- About 10% to 20% of mesotheliomas are **sarcomatoid** (**fibrous**).
- **Mixed (biphasic)** mesotheliomas have both epithelioid and sarcomatoid areas. They make up about 30% to 40% of mesotheliomas.

Benign tumors of the mesothelium

Benign (non-cancerous) tumors can also start in the mesothelium. These tumors are typically removed by surgery, and there is often no need for additional treatment.

Localized fibrous tumor of the pleura

This type of benign tumor can form in the pleura surrounding the lungs. It used to be called *benign fibrous mesothelioma*, but doctors now know that this tumor actually starts from tissue under the mesothelium and not from mesothelial cells. This disease is usually benign, but about 1 in 10 are cancerous. A similar condition that starts in the peritoneum is called *solitary fibrous tumor of the peritoneum*.

Adenomatoid mesothelioma

This benign tumor can develop in the mesothelium of certain reproductive organs. In men, it often starts in the epididymis (ducts that carry sperm cells out of the testicle). In women, this tumor may begin in the fallopian tubes (tubes that carry eggs from the ovaries to the uterus).

Benign cystic mesothelioma

This rare non-cancerous tumor often begins in the peritoneum.

Only malignant mesothelioma will be discussed further in this document.

What are the key statistics about malignant mesothelioma?

Mesothelioma is fairly rare. About 3,000 new cases of mesothelioma are diagnosed each year in the United States.

The rate of mesotheliomas in the United States increased from the 1970s to the early 1990s, but since then it has leveled off and even gone down slightly. These changes have largely been seen in men, and are probably related to changes in workplace exposures to asbestos (see "What are the risk factors for malignant mesothelioma?"). The rate of mesothelioma is lower in women and has been fairly steady for some time. In many other countries, the rate of mesothelioma is still increasing.

Mesothelioma is more common in whites and Hispanics/Latinos than in African Americans or Asian Americans.

Information on survival rates for mesothelioma can be found in the section, "Survival statistics for mesothelioma."

What are the risk factors for malignant mesothelioma?

A risk factor is anything that affects your chance of getting a disease such as cancer. Different cancers have different risk factors. Some risk factors, like smoking, can be changed. Others, like a person's age or family history, can't be changed.

But risk factors don't tell us everything. Having a known risk factor, or even several risk factors, does not mean that you will get the disease. And some people who get the disease may have few or no known risk factors.

Researchers have found some factors that increase a person's risk of mesothelioma.

Asbestos

The main risk factor for developing mesothelioma is exposure to asbestos. In fact, up to 3 of every 4 cases of mesothelioma have been linked to asbestos exposure. Most often, people are exposed in the workplace.

Asbestos is a group of minerals that occur naturally as bundles of fibers. These fibers, found in soil and rocks in many parts of the world, are made of silicon, oxygen, and other elements.

When asbestos fibers in the air are inhaled, they tend to stick to mucus in the throat, trachea (windpipe), or bronchi (large breathing tubes of the lungs). Fibers that stay in the lungs can travel to the ends of the small airways and penetrate the pleural lining of the

lung and chest wall. These fibers may then injure the cells of the pleura, and eventually cause mesothelioma.

Asbestos fibers can also damage cells of the lung and result in *asbestosis* (scar tissue in the lung) and/or lung cancer. Indeed, asbestosis, mesothelioma, and lung cancer are the 3 most frequent causes of death and disease among people with heavy asbestos exposure.

Peritoneal mesothelioma, which forms in the abdomen, may result from coughing up and swallowing inhaled asbestos fibers.

Many people are exposed to very low levels of naturally occurring asbestos in outdoor air in dust that comes from rocks and soil containing asbestos. This is more likely to happen in areas where rocks have higher asbestos content. In some areas, asbestos may be found in the water supply as well as in the air.

In the past, asbestos was used in many products because of its heat and fire-resistant properties. The link between asbestos and mesothelioma is well known, so its use in the United States has gone down dramatically. Most use stopped after 1989, but it is still used in some products.

Still, millions of Americans may already have been exposed to asbestos. People at risk for asbestos exposure in the workplace include some miners, factory workers, insulation manufacturers and installers, railroad and automotive workers, ship builders, gas mask manufacturers, and construction workers. Family members of people exposed to asbestos at work can also have an increased risk of developing mesothelioma because the workers can carry home asbestos fibers on their clothes. The rate of mesothelioma in men appears to be dropping, probably because there is now much less direct exposure to asbestos in the workplace.

Asbestos was also used to insulate many older homes, as well as commercial and public buildings around the country, including some schools. Because these particles are contained within the building materials, they are not likely to be found in the air in large numbers. The risk of exposure is likely to be very low unless the particles somehow escape into the air, such as when building materials begin to decompose over time, or during remodeling or removal.

The risk of developing mesothelioma is related to how much asbestos a person was exposed to and how long this exposure lasted. People exposed at an early age, for a long period of time, and at higher levels are more likely to develop this cancer. Mesotheliomas take a long time to develop. The time between first exposure to asbestos and diagnosis of mesothelioma is usually between 20 and 50 years. Unfortunately, the risk of mesothelioma does not go down over time after the exposure to asbestos stops. The risk appears to be lifelong.

For more detailed information on asbestos, see our document, Asbestos.

Zeolites

Zeolites are minerals that are chemically related to asbestos. An example is erionite, which is common in the rocks and soil in parts of Turkey. High mesothelioma rates in these areas are believed to be caused by exposure to this mineral.

Radiation

There have been a few published reports of mesotheliomas that developed after people were exposed to high doses of radiation to the chest or abdomen as treatment for another cancer. Although the risk of mesothelioma is increased in patients who have been treated with radiation, this cancer still only occurs rarely in these patients.

There have also been reports linking mesothelioma to injections of thorium dioxide (Thorotrast). This radioactive material was used by doctors for certain x-ray tests until the 1950s. Thorotrast was found to cause cancers, so it has not been used for many years.

SV40 virus

Some studies have raised the possibility that infection with simian virus 40 (SV40) might increase the risk of developing mesothelioma. Some injectable polio vaccines given between 1955 and 1963 were contaminated with SV40. As many as 30 million people in the United States may have been exposed to this virus.

Some lab studies have suggested that SV40 infection might cause mesothelioma. For example, infecting some lab animals like hamsters with SV40 causes mesotheliomas to develop. Researchers also have noticed that SV40 can cause mouse cells grown in lab dishes to become cancerous, and that asbestos increases the cancer-causing effect of SV40 on these cells. Other researchers have found SV40 DNA in some biopsy specimens of human mesotheliomas. But fragments of SV40 DNA can also be found in some non-cancerous human tissues.

So far, the largest studies looking at this issue in humans have not found an increased risk for mesothelioma or other cancers among people who received the contaminated vaccines as children. But the peak age range for diagnosis of mesothelioma is 50 to 70 years. Some researchers have pointed out that this issue may remain unresolved until more of the people accidentally exposed to SV40 between 1955 and 1963 reach that age range.

Most experts have concluded that at this time we still don't know if SV40 is responsible for some mesotheliomas. This important topic is still being researched.

Age

The risk of mesothelioma increases with age. It is rare in people under age 45. About 2 out of 3 people with mesothelioma of the chest are 65 or older.

Gender

The disease is much more common in men than in women. This is probably because men have been more likely to work in jobs with heavy exposure to asbestos.

Do we know what causes malignant mesothelioma?

Cancers, including mesotheliomas, occur when cells in the body suffer damage to their DNA. DNA is the chemical in each of our cells that makes up our *genes* - the instructions for how our cells function. We usually look like our parents because they are the source of our DNA. But DNA affects more than how we look. Some genes control when cells in the body grow and divide into new cells. Changes in these genes may cause cells to grow out of control, which can lead to cancer.

Asbestos exposure is the main cause of mesothelioma. After these fibers are breathed in, they travel to the ends of small air passages and reach the pleura, where they can damage mesothelial cells. This leads to inflammation and scarring. This may damage cells' DNA and cause changes that result in uncontrolled cell growth. If swallowed, these fibers can reach the abdominal cavity where they have a role in causing peritoneal mesothelioma.

But most people exposed to asbestos, even in large amounts, do not get mesothelioma. Other factors, such as a person's genes, may make them more likely to develop mesothelioma when exposed to asbestos. For example, researchers have found that some people who seem to be at high risk have changes in *BAP1*, a gene that normally helps keep cell growth under control. Other genes are probably important as well.

Radiation treatments for other cancers have been linked to mesothelioma in some studies. Radiation can damage the cells' DNA, leading to out-of-control cell growth.

It is still not known if infection with the SV40 virus increases the risk of mesothelioma, or exactly how it might do so. In lab studies, researchers have found that the virus can affect certain genes that have been linked with cancer, but further research in this area is needed.

Researchers now understand many of the factors that increase a person's risk of mesothelioma, but it's still not clear exactly how these factors cause the gene changes that lead to cancer. This is an active area of research.

Can malignant mesothelioma be prevented?

Being exposed to asbestos is by far the biggest risk factor for mesothelioma, so the best way to reduce your risk is to limit your exposure to asbestos in homes, in public buildings, and at work.

People who might be exposed to high levels of asbestos at work include some miners, factory workers, insulation manufacturers and installers, railroad and automotive

workers, ship builders, gas mask manufacturers, and construction workers. If there is a chance of on-the-job exposure, such as during the renovation of old buildings, then you should use all protective equipment and safety procedures designed for working around asbestos.

Older homes may have insulation containing asbestos or other materials. A knowledgeable expert can check your home to find out if there is any asbestos and whether it poses any risk of exposure. This may mean testing the air for asbestos levels. Just because asbestos exists in a home does not necessarily mean that it needs to be removed. As long as the material is not damaged or disturbed, for example by drilling or remodeling, the fibers will not be released into the air. If asbestos needs to be removed from your home, you should hire a qualified contractor to do this to avoid contaminating your home or causing any exposure to your family or to the workers. You should not attempt to remove asbestos-containing material yourself.

Asbestos can also be found in some commercial and public buildings (including some schools), where the same basic principles apply. Intact, undisturbed materials containing asbestos generally do not pose a health risk. They may pose a risk if they are damaged, are disturbed in some way, or deteriorate over time and release asbestos fibers into the air. By federal law, all schools are required to inspect materials with asbestos regularly and to have a plan in place for managing them.

Can malignant mesothelioma be found early?

Mesothelioma is uncommon, and there are no widely recommended screening tests for this cancer in people who are not at increased risk. (Screening is testing for cancer in people who don't have any symptoms.)

For people with known exposure to asbestos, some doctors recommend imaging tests such as chest x-rays or computed tomography (CT) scans to look for changes in the lungs that might be signs of mesothelioma or lung cancer. But it is not clear how useful these tests are in finding mesotheliomas early.

In recent years, doctors have found that people with mesothelioma often have high levels of certain substances in their blood, including *osteopontin* and *soluble mesothelin-related peptides* (SMRPs). Blood tests for these substances may one day be useful in finding mesotheliomas early, and in monitoring the course of the disease in people who have mesothelioma.

Most mesotheliomas are found when a person goes to a doctor because of symptoms. People who have been exposed to asbestos should know the possible signs and symptoms of mesothelioma (discussed in the next section). Many of these symptoms are more likely to be caused by something other than mesothelioma, but it's important to report any new symptoms to your doctor right away so that the cause can be found and treated, if needed.

Signs and symptoms of mesothelioma

Early symptoms of mesothelioma are more often caused by other things, so at first people may ignore them or mistake them for common, minor ailments. Most people with mesothelioma have symptoms for at least a few months before they are diagnosed.

Symptoms of pleural mesothelioma (mesothelioma of the chest) can include:

- Pain in the lower back or at the side of the chest
- Shortness of breath
- Fluid in the area around the lung
- Cough
- Fever
- Excessive sweating
- Fatigue
- Weight loss (without trying)
- Trouble swallowing (feeling like food gets stuck)
- Hoarseness
- Swelling of the face and arms

Symptoms of peritoneal mesothelioma can include:

- Abdominal (belly) pain
- Swelling or fluid in the abdomen
- Weight loss (without trying)
- Nausea and vomiting

The symptoms and signs above may be caused by mesothelioma, but more often they are caused by other conditions. Still, if you have any of these problems (especially if you have been exposed to asbestos), it's important to see your doctor right away so the cause can be found and treated, if needed.

How is malignant mesothelioma diagnosed?

Mesothelioma is most often diagnosed after a person goes to a doctor because of symptoms. If there is a reason to suspect you might have mesothelioma, your doctor will use one or more tests to find out. Symptoms might suggest that the problem could be mesothelioma, but tests will be needed to confirm the diagnosis.

Medical history and physical exam

If you have any signs or symptoms that suggest you might have mesothelioma, your doctor will want to take a complete medical history to learn about your symptoms and possible risk factors, especially asbestos exposure. You will also be asked about your general health.

A physical exam can provide information about possible signs of mesothelioma and other health problems. Pleural mesothelioma can cause fluid to build up around the lungs in the chest (called a *pleural effusion*). In cases of peritoneal mesothelioma, fluid can build up in the abdomen (called *ascites*). In pericardial mesothelioma, fluid builds up in the sac around the heart (called a *pericardial effusion*). Rarely, mesothelioma can develop in the groin and look like a hernia. All of these might be found during a physical exam, such as when the doctor listens to these areas with a stethoscope.

If mesothelioma is a possibility, tests will be needed to make sure. These might include imaging tests, blood tests, and other procedures.

Imaging tests

Imaging tests use x-rays, radioactive particles, or magnetic fields to create pictures of the inside of your body. Imaging tests may be done for a number of reasons, including to help find a suspicious area that might be cancerous, to learn how far cancer may have spread, and to help determine if treatment has been effective.

Chest x-ray

This is often the first test done if someone has symptoms such as a constant cough or shortness of breath. It may show an abnormal thickening of the pleura, calcium deposits on the pleura, fluid in the space between the lungs and the chest wall, or changes in the lungs themselves as a result of asbestos exposure. These findings could suggest a mesothelioma.

Computed tomography (CT) scan

The CT scan is an x-ray test that produces detailed cross-sectional images of your body. Instead of taking one picture, like a regular x-ray, a CT scanner takes many pictures as it rotates around you while you are lying on a narrow table. A computer then combines these into images of slices of the body. Unlike a regular x-ray, a CT scan creates detailed images of the soft tissues in the body.

A CT scanner has been described as a large donut, with a narrow table in the middle opening. You will need to lie still on the table while the scan is being done. CT scans take longer than regular x-rays, and you might feel a bit confined by the ring while the pictures are being taken.

CT scans are often used to help look for mesothelioma and to determine the exact location of the cancer. They can also help stage the cancer (determine the extent of its spread). For example, they can show whether the cancer has spread to other organs. This can help to determine if surgery might be a treatment option. Finally, CT scans can be used to learn whether treatment such as chemotherapy has been helpful in shrinking or slowing the growth of the cancer.

Before the test, you might have to drink 1 to 2 pints of a liquid called *oral contrast*. This helps outline the intestine so that certain areas are not mistaken for tumors. You might also receive an IV (intravenous) line through which a different kind of contrast dye (IV contrast) is injected. This helps better outline structures in your body. The injection can cause some flushing (redness and warm feeling). Some people are allergic and get hives or, rarely, more serious reactions like trouble breathing and low blood pressure. Be sure to tell the doctor if you have any allergies or have ever had a reaction to any contrast material used for x-rays.

Echocardiogram

This test uses sound waves to look at the heart. It may be done if your doctor suspects that you have fluid around your heart (a pericardial effusion). It can also tell how well the heart is working. For the most common version of this test, you lie on a table while a technician moves an instrument called a *transducer* over the skin on your chest. A gel is often put on the skin first.

Positron emission tomography (PET) scan

For a PET scan, a radioactive substance (usually a type of sugar related to glucose, known as *FDG*) is injected into the blood. The amount of radioactivity used is very low. Because cancer cells in the body grow quickly, they absorb more of the sugar than most other cells. After waiting about an hour, you lie on a table in the PET scanner for about 30 minutes while a special camera creates a picture of areas of radioactivity in the body.

The picture from a PET scan is not finely detailed like a CT or MRI scan, but it can provide helpful information about whether abnormal areas seen on these tests are likely to be cancerous or not. For example, it can give the doctor a better idea of whether a thickening of the pleura or peritoneum seen on a CT scan is more likely cancer or merely scar tissue. If you have been diagnosed with cancer, your doctor may use this test to see if the cancer has spread to lymph nodes or other parts of the body. A PET scan can also be useful if your doctor thinks the cancer may have spread but doesn't know where.

Some machines can do both a PET and CT scan at the same time (PET/CT scan). This lets the doctor compare areas of higher radioactivity on the PET scan with the more detailed appearance of that area on the CT.

Magnetic resonance imaging (MRI) scan

Like CT scans, MRI scans make detailed images of soft tissues in the body. But MRI scans use radio waves and strong magnets instead of x-rays. The energy from the radio waves is absorbed and then released in a pattern formed by the type of body tissue and by certain diseases. A computer translates the pattern into very detailed images of parts of the body. A contrast material called *gadolinium* is often injected into a vein before the scan to better show details. This contrast is different than the one used for CT scans, so being allergic to one doesn't mean you are allergic to the other.

MRI scans can sometimes help show the exact location and extent of a tumor since they provide very detailed images of soft tissues. For mesotheliomas, they may be useful in looking at the diaphragm (the thin band of muscle below the lungs that helps us breathe), a possible site of cancer spread.

MRI scans take longer than CT scans – often up to an hour. You may have to lie inside a narrow tube, which can upset people with a fear of enclosed spaces. Special, more open MRI machines may be an option in some cases. The MRI machine makes buzzing and clicking noises that you might find disturbing. Some places will give you earplugs to help block this out.

More information on radiation therapy can be found in the Radiation section of our website, or in our document *Understanding Radiation Therapy: A Guide for Patients and Families*

Blood tests

Blood levels of certain substances are often higher in people with mesothelioma:

- Osteopontin
- Soluble mesothelin-related peptides (SMRPs), detected with the MesoMark® test

Blood tests for these substances are not used to diagnose mesothelioma, but high levels may make the diagnosis more likely.

If you have been diagnosed with mesothelioma, other blood tests will be done to check your blood cell counts and levels of certain chemicals in the blood. These tests can give the doctor an idea of how extensive the disease may be, and how well organs such as the liver and kidneys are working.

Tests of fluid and tissue samples

Symptoms and test results may strongly suggest that a person has mesothelioma, but the actual diagnosis is made by removing cells from an abnormal area and looking at them under a microscope. This is known as a *biopsy*. It may be done in different ways, depending on the situation.

Removing fluid for testing

If mesothelioma might be causing a buildup of fluid in the body, a sample of this fluid can be removed by inserting a thin, hollow needle through the skin and into the fluid to remove it. Numbing medicine is used on the skin before the needle is inserted. This may be done in a doctor's office or in the hospital. Sometimes ultrasound (or an echocardiogram) is used to guide the needle (these use sound waves to take pictures of parts of the body).

This procedure has different names depending on where the fluid is:

- Thoracentesis removes fluid from the chest.
- Paracentesis removes fluid from the abdomen.
- **Pericardiocentesis** removes fluid from the sac around the heart.

The fluid is then tested to check its chemical makeup and is looked at under a microscope to see if it contains cancer cells. If cancer cells are found, special tests might be able to tell whether the cancer is a mesothelioma, a lung cancer, or another type of cancer.

Even if no cancer cells are found in the fluid, cancer may still be present. In many cases, doctors need to get an actual sample of the mesothelium (the pleura, peritoneum, or pericardium) to determine if a person has mesothelioma.

Needle biopsies

Suspected tumors in the chest are sometimes sampled by needle biopsy. A long, hollow needle is passed through the skin in the chest between the ribs and into the pleura. Imaging tests such as CT scans are used to guide the needle into the tumor so that small samples can be removed to be looked at under the microscope. This is often done using just numbing medicine.

Needle biopsy can also be used to get samples of the lymph nodes in the space between the lungs to see if the cancer has spread there (see "Endobronchial ultrasound needle biopsy").

Needle biopsies do not require a surgical incision or overnight hospital stay. But the downside is that sometimes the samples removed are not big enough to make an accurate diagnosis. This is especially true for mesothelioma. A more invasive biopsy method may be needed.

There is a slight chance that the needle could put a small hole in the lung during the biopsy. This can cause air to build up in the space between the lung and the chest wall (known as a *pneumothorax*). A small pneumothorax may cause no symptoms. It may only be seen on an x-ray done after the biopsy, and it will often go away on its own. But a larger pneumothorax can make part of a lung collapse and might need to be treated. The treatment is placement of a small tube (a catheter) through the skin and into the space

between the lungs. The tube is used to suck the air out in order to re-expand the lung and is left in place for a short time.

Endoscopic biopsies

An endoscope is a thin, tube-like instrument used to look inside the body. It has a light and a lens (or tiny video camera) on the end for viewing and often has a tool to remove tissue samples. Endoscopes have different names depending on the part of the body where they're used. Endoscopic biopsy is commonly used to diagnose mesothelioma.

Thoracoscopy: This procedure uses an endoscope called a *thoracoscope* to look at areas inside the chest. It can be used to look at the pleura and take tissue samples for biopsies. Thoracoscopy is done in the operating room while you are under general anesthesia (in a deep sleep). The doctor inserts the thoracoscope through one or more small cuts made in the chest wall to look at the space between the lungs and the chest wall. This lets the doctor see possible areas of cancer and remove small pieces of tissue to look at under the microscope. The doctor can also sample lymph nodes and fluid and see if a tumor is growing into nearby tissues or organs. Thoracoscopy can also be used as part of a procedure to keep fluid from building up in the chest. This is called *pleurodesis* and is discussed in the section "Palliative procedures used for malignant mesothelioma."

Laparoscopy: For this test, the doctor uses an endoscope called a *laparoscope* to look inside the abdomen and biopsy any peritoneal tumors. This is done in the operating room while you are under general anesthesia (in a deep sleep). The laparoscope is inserted into the abdomen through small cuts on the front of the abdomen.

Mediastinoscopy: If imaging tests such as a CT scan suggest that the cancer has spread to the lymph nodes between the lungs, the doctor may want to remove some of them to see if they really contain cancer. This area is the mediastinum, and looking at it with an endoscope is called *mediastinoscopy*. This is also done in an operating room while you are under general anesthesia (in a deep sleep).

A small cut is made in the front of the neck above the breastbone (sternum) and a thin, hollow, lighted tube (called a *mediastinoscope*) is inserted behind the sternum. Special instruments can be passed through this tube to take tissue samples from the lymph nodes along the windpipe and the major bronchial tube areas.

Cancers in the lung often spread to lymph nodes, but mesotheliomas do this less often. Testing the lymph nodes can help show whether a cancer is still localized or if it has started to spread, which might affect treatment options. It can also sometimes help distinguish lung cancer from mesothelioma.

Bronchoscopy: This test does not require any cuts in the skin. It uses a bronchoscope – a long, thin, flexible, fiber-optic tube that is placed down the throat and into the lungs to look at the lining of the main airways. This procedure is done while you are asleep or sedated. If a tumor is found, the doctor can take a small sample of the tumor through the tube. *endobronchial ultrasound needle biopsy*. For this, a bronchoscope with an ultrasound device at its tip is passed down into the windpipe. The ultrasound device uses

sound waves to let the doctor see the nearby lymph nodes. A hollow needle is then passed down the bronchoscope and through the airway wall into the nodes to take biopsy samples. This procedure may be done with either general anesthesia (you are asleep), or with numbing medicine (local anesthesia) and light sedation.

Open surgical biopsy

Sometimes, endoscopic biopsies aren't enough to make a diagnosis and more invasive procedures are needed. By making an incision in the chest (thoracotomy) or an incision in the abdomen (laparotomy) the surgeon can remove a larger sample of tumor or, sometimes, remove the entire tumor.

Testing the samples in the lab

No matter how they were obtained, all biopsy and fluid samples are sent to the pathology lab. There, a doctor will look at them under a microscope and test them to find out if they contain cancer cells (and if so, what type of cancer it is).

It is often hard to diagnose mesothelioma by looking at cells from fluid around the lungs, abdomen, or heart. It can even be hard to diagnose mesothelioma with tissue from small needle biopsies. Under the microscope, mesothelioma can often look like other types of cancer. For example, pleural mesothelioma may resemble some types of lung cancer, and peritoneal mesothelioma in women may look like some cancers of the ovaries.

For this reason, special lab tests are often done to help tell mesothelioma from some other cancers.

Immunohistochemistry: This test looks for different proteins on the surface of the cells or inside them. It can be used to tell if the cancer is a mesothelioma or a lung cancer, which can sometimes appear to start in the inner lining of the chest.

DNA microarray analysis: This is a newer test that actually looks at patterns of genes in the cancer cells. Mesothelioma cells have different gene patterns than other cancer cells.

Electron microscopy: This can sometimes help diagnose mesothelioma. The electron microscope can magnify samples many more times than a normal light microscope. This more powerful microscope makes it possible to see the small parts of the cancer cells that can distinguish mesothelioma from other types of cancer.

For more about these and other tests that are done on tissue samples, you can read our document *Testing Biopsy and Cytology Specimens for Cancer*.

If mesothelioma is diagnosed, the doctor will also determine what type of mesothelioma it is, based on the patterns of cells seen in the microscope. Most mesotheliomas are classified as either epithelioid, sarcomatoid, or mixed/biphasic.

Pulmonary function tests

If mesothelioma has been diagnosed, pulmonary function tests (PFTs) may be done to see how well your lungs are working. This is especially important if surgery may be an option to treat the cancer. Surgery often requires removing part or all of a lung, so it's important to know how well the lungs are working to start with. These tests can give the surgeon an idea of whether surgery may be an option, and if so, how much lung can safely be removed.

There are a few different types of PFTs, but they all basically have you breathe in and out through a tube that is connected to a machine that measures your lung function.

How is malignant mesothelioma staged?

Staging is the process of finding out how far a cancer has spread. Your treatment and prognosis (outlook) depend, to a large extent, on the cancer's stage.

Staging is based on the results of physical exams, biopsies, and imaging tests (CT scan, PET scan, etc.), which are described in the section, "How is malignant mesothelioma diagnosed?"

A staging system is a formal way for doctors and nurses to sum up the extent of a cancer. Pleural mesothelioma, the most common type, is the only mesothelioma for which a formal staging system exists.

The TNM staging system

The system most often used to describe the growth and spread of pleural mesothelioma is the American Joint Committee on Cancer (AJCC) **TNM** staging system. The TNM system is based on 3 key pieces of information:

- T sums up the extent of spread of the main (primary) tumor.
- N describes how much the cancer has spread to nearby (regional) lymph **nodes**. Lymph nodes are small bean-shaped collections of immune system cells to which cancers often spread first.
- M indicates whether the cancer has spread (metastasized) to other organs of the body. (The most common sites are the pleura on the other side of the body, the lungs, and the peritoneum.)


Numbers or letters appear after T, N, and M to provide more details about each of these factors. Higher numbers mean the cancer is more advanced.

T groups

TX: The main tumor can't be assessed for some reason.

- **T0:** There is no evidence of a main tumor (the cancer is found elsewhere instead).
- **T1:** Mesothelioma involves the pleura lining the chest wall on one side of the chest. It may or may not also affect the pleura lining the diaphragm (the breathing muscle) or the mediastinum (the space between the lungs). It may also have spread to the pleura covering the lung.
- **T2:** Mesothelioma involves the pleura lining the chest wall on one side of the chest. It also involves the pleura coating the diaphragm, the mediastinum, and the lung. It also has grown into at least one of the following:
 - The diaphragm
 - The lung itself
- **T3:** The mesothelioma has grown further but may still possibly be removed with surgery. The tumor involves the pleura lining the chest wall on one side of the chest. It also involves the pleura coating the lung, the diaphragm, and the mediastinum. It also has grown into at least one of the following:
 - The first layer of the chest wall (called the *endothoracic fascia*)
 - The fatty tissue in the mediastinum
 - A single place in the deeper layers of the chest wall
 - The surface of the pericardium (outer covering layer of the heart)
- **T4:** The mesothelioma has grown too far to be removed completely with surgery. The tumor has grown into the pleura lining the chest wall on one side of the chest. It also is in the pleura coating the lung, diaphragm, and mediastinum on the same side. It also has grown into at least one of the following:
 - More than one place in the deeper layers of the chest wall, including the muscle or ribs
 - Through the diaphragm and into the peritoneum
 - Any organ in the mediastinum (esophagus, trachea, thymus, blood vessels)
 - The spine
 - Across to the pleura on the other side of the chest
 - Through the heart lining (pericardium) or into the heart itself

N groups



NX: The nearby lymph nodes can't be assessed.

N0: No spread to nearby lymph nodes.

N1: Spread to lymph nodes within the lung and/or around the area where the bronchus enters the lung (called *hilar* or *bronchial* lymph nodes) on the same side as the main tumor.

N2: Spread to other lymph nodes on the same side as the main tumor, such as the *subcarinal* (around the point where the windpipe branches into the left and right bronchi) and the *mediastinal* lymph nodes. Also includes spread to the lymph nodes in the space just behind the breastbone (called *internal mammary* lymph nodes) and those near the diaphragm (called *peridiaphragmatic*).

N3: Spread to lymph nodes near the collarbone on either side (*supraclavicular* lymph nodes), and/or spread to hilar or mediastinal lymph nodes on the side opposite the main tumor.

M groups

M0: No spread to distant organs or areas.

M1: The cancer has spread to distant sites. This can be to distant lymph nodes or to other organs.

Stage grouping for pleural mesothelioma

Once the T, N, and M categories have been assigned, this information is combined in a process called stage grouping to assign an overall stage of I, II, III, or IV. The stages identify tumors that have a similar prognosis and thus are treated in a similar way. Patients with lower stage numbers tend to have a better prognosis.

Stage I

T1, N0, M0: Mesothelioma has grown into the pleura lining the chest wall on one side of the chest. It might or might not also affect the pleura lining the diaphragm (the breathing muscle) or the mediastinum (the space between the lungs). It may also have spread to the pleura covering the lung (T1). It has not spread to the lymph nodes (N0) or to distant sites (M0).

Stage II

T2, N0, M0: Mesothelioma has grown into the pleura lining the chest wall on one side of the chest. It also is in the pleura coating the diaphragm, the mediastinum, and the lung. The cancer has also grown into the diaphragm or the lung itself (T2). It has not spread to the lymph nodes (N0) or to distant sites (M0).

Stage III

Either of the following:

T1 or T2, N1 or N2, M0: Mesothelioma has grown into the pleura lining the chest on one side, and might or might not have grown into the pleura lining the lung, the diaphragm, or the mediastinum. It might also have grown into the muscle of the diaphragm or the lung itself (T1 or T2). It has spread to lymph nodes in the chest on the same side as the main tumor (N1 or N2). It has not spread to distant sites (M0).

OR

T3, N0 to N2, M0: Mesothelioma is in the pleura lining the chest on one side, and has grown into the first layer of the chest wall, the fatty tissue in the mediastinum, a single place in the deeper layers of the chest wall, or the outer covering layer of the heart. It might or might not have spread to lymph nodes in the chest on the same side as the tumor but has not spread to lymph nodes near the collarbone or on the opposite side of the chest (N0, N1, or N2). It has not spread to distant sites (M0).

Stage IV

Any of the following:

T4, any N, M0: Mesothelioma has grown into the pleura lining the chest on one side and has grown into more than one place in the deeper layers of the chest wall (including the muscle or ribs), through the diaphragm and into the peritoneum, into any organ in the mediastinum, into the spine, across to the pleura on the other side of the chest, and/or through the heart lining or into the heart itself. It might or might not have spread to lymph nodes (any N). It has not spread to distant sites (M0).

Any T, N3, M0: The tumor may or may not have grown into nearby tissues (any T). It has spread to lymph nodes near the collarbone on either side and/or to hilar or mediastinal lymph nodes on the side opposite the main tumor (N3). It has not spread to distant sites (M0).

OR

Any T, any N, M1: The mesothelioma might or might not have grown into nearby tissues (any T). It might or might not have spread to the lymph nodes (any N). It has spread to distant sites (M0).

Resectable versus unresectable cancer

The TNM system divides mesotheliomas into several stages that help give doctors an idea about a person's prognosis (outlook). But for treatment purposes, doctors often use a simpler system based on whether these cancers are likely to be resectable (where all visible tumor can be removed by surgery) or unresectable.

In general terms, most stage I, II, and III mesotheliomas are potentially resectable, but there are exceptions. Whether or not the cancer can be removed depends not only on how far the tumor has grown into nearby tissues, but also on its subtype (most doctors believe only epithelioid and mixed/biphasic tumors are potentially resectable), where it is located, and if the patient is healthy enough to have surgery.

Even for resectable mesotheliomas, in most cases cancer cells that cannot be seen are left behind after surgery. For this reason, many doctors use other forms of treatment (radiation therapy and/or chemotherapy) along with surgery when possible.

Other prognostic factors

Stage is an important factor in predicting a patient's prognosis (outlook), but other factors also play a role. Some factors linked to longer survival times include:

- Good performance status (being able to carry out normal tasks of daily life)
- Younger age
- Epithelioid subtype
- Not having chest pain
- No significant weight loss
- Normal levels of a substance in the blood called LDH
- Normal red blood cell counts, white blood cell counts, and blood platelet counts

Survival statistics for mesothelioma

Survival rates are often used by doctors as a standard way of discussing a person's prognosis (outlook). Some patients want to know the survival statistics for people in similar situations, while others may not find the numbers helpful, or even don't want to know them. If you don't want to read about the survival statistics for mesothelioma, stop here and skip to the next section.

To get survival rates, doctors have to look at people who were treated at least several years ago. Although the numbers below are among the most current we have available, improvements in treatment since then could result in a more favorable outcome for people now being diagnosed with mesothelioma.

Survival rates are often based on previous outcomes of large numbers of people who had the disease, but they cannot be used to predict what will happen in any particular person's case. Knowing the type and the stage of a person's cancer is important in estimating their outcome. But many other factors can affect survival, such as a person's age and overall health, the treatment received, and how well the cancer responds to treatment. Even taking these other factors into account, survival rates are at best rough estimates. Your doctor can tell you if the numbers below apply, as he or she is familiar with your situation.

Mesothelioma is a serious disease. By the time the symptoms appear and cancer is diagnosed, the disease is often advanced. Regardless of the extent of the cancer, mesothelioma can be very hard to treat.

Relative 5-year survival takes the proportion of people with the cancer that have survived 5 years (the 5 year survival) and compares it to the survival expected in a similar group of people without the cancer. This helps adjust for deaths from causes other than the cancer. Based on data from the National Cancer Institute's SEER program, the relative 5-year survival rate for mesothelioma is between 5% and 10%. People diagnosed at a younger age tend to survive longer.

The numbers in the table below are from a large international study that looked at the median survival time of patients with pleural mesothelioma who were treated with surgery to cure the cancer. The numbers include the relative 5-year survival rate and median survival. Median survival is the length of time it took for half the people in a certain group (like those with a certain type and stage of cancer) to die. It is kind of like an average – half the patients in the group live longer than that and half the patients don't.

Stage	Median Survival
Ι	21 months
II	19 months

III	16 months
IV	12 months

As a general rule, survival times are likely to be longer for people with mesotheliomas that can be operated on than for those with cancers that have spread too far to be removed. Other prognostic factors, such as those listed in the previous section, also affect survival.

How is malignant mesothelioma treated?

This information represents the views of the doctors and nurses serving on the American Cancer Society's Cancer Information Database Editorial Board. These views are based on their interpretation of studies published in medical journals, as well as their own professional experience.

The treatment information in this document is not official policy of the Society and is not intended as medical advice to replace the expertise and judgment of your cancer care team. It is intended to help you and your family make informed decisions, together with your doctor.

Your doctor may have reasons for suggesting a treatment plan different from these general treatment options. Don't hesitate to ask him or her questions about your treatment options. If you have mesothelioma, your cancer care team will recommend one or more treatment options for you to consider. This is an important decision and you should take time to think about all of your choices.

Making treatment decisions

After the cancer is found and staged, your cancer care team will discuss your treatment options with you. The main factors in selecting treatment for mesotheliomas are the location and extent of the tumor, whether it has spread to lymph nodes or other organs, and your health and personal preferences. Based on these factors, your treatment options may include:

- Surgery
- Radiation therapy
- Chemotherapy

More than one of these treatments may be used in many cases.

Because mesothelioma is a rare cancer, it has been hard for doctors to compare the value of different treatments. Only a few large clinical trials of treatments for mesothelioma have been done. In addition, many doctors have little or no experience treating this disease. They usually refer patients with this cancer to specialists who treat large numbers of mesothelioma patients at major medical centers.

You may have different types of doctors on your treatment team, depending on the stage of your cancer and your treatment options. These doctors may include:

- A thoracic surgeon: a doctor who treats diseases of the lungs and chest with surgery.
- A surgical oncologist: a doctor who treats cancer with surgery
- A radiation oncologist: a doctor who treats cancer with radiation therapy.
- A medical oncologist: a doctor who treats cancer with medicines such as chemotherapy.
- A pulmonologist: a doctor who specializes in medical treatment of diseases of the lungs.

Many other specialists may be involved in your care as well, including nurse practitioners, nurses, respiratory therapists, social workers, and other health professionals.

Mesothelioma is often hard to treat because it typically does not grow as a single tumor mass. It tends to spread along nearby surfaces, nerves, and blood vessels. This often makes it very hard to get rid of it completely with surgery and/or radiation.

Before deciding on a treatment plan, it's very important to have an idea of its likely benefits and possible risks. You will probably have many questions about the treatment options suggested. Mesotheliomas are rare, so if time permits it is often a good idea to get a second opinion from a doctor who has a lot of experience in treating people with these cancers. A second opinion can give you more information and help you feel more confident about the treatment plan you choose.

The next few sections describe the types of treatments used for mesotheliomas. This is followed by a discussion of the most common treatments used, based on the extent of the disease.

Surgery for malignant mesothelioma

Surgery for mesothelioma may be done for 1 of 2 reasons:

- To try to cure the cancer
- To relieve (palliate) pain and other symptoms caused by the tumor

Surgery to try to cure the cancer is known as *potentially curative surgery*. This type of surgery may be an option if you are in otherwise good health and the cancer has not spread too far to be removed completely. Unfortunately, even when the surgeon can remove all of the cancer that can be seen, some cancer cells are often left behind. These cells can grow and divide, causing the cancer to come back after surgery. Because of this, not all doctors agree on the exact role of surgery. In most cases it is not likely to cure you but may extend your life. Still, potentially curative surgery is being done in some major cancer centers, and a small number of patients who have had the surgery have had long remissions of their disease.

Palliative surgery may be an option if the tumor has already spread beyond where it started and is difficult to remove completely, or if you are too ill for a more extensive

operation. The goal of this surgery is to relieve or prevent symptoms, as opposed to trying to cure the cancer.

Surgery for pleural mesothelioma

Surgery for pleural mesothelioma may be done either to help prevent or relieve symptoms or to try to remove all of the cancer. Unfortunately, these tumors often have spread too far to be removed completely. Sometimes, the surgeon may not be able to tell the full extent of the cancer – and therefore which type of surgery might be best – until the operation has started.

Extrapleural pneumonectomy (EPP): This is an extensive operation but it may offer the best chance to remove all of the cancer for many patients. It is most often used when the surgeon thinks a cure is possible – typically in patients with resectable epithelioid mesothelioma that has not spread to the lymph nodes.

This operation removes the lung on the side of the cancer along with the pleura lining the chest wall on that side, the diaphragm on that side, the pericardium (the sac around the heart), and nearby lymph nodes. The diaphragm and the pericardium are then reconstructed with man-made materials.

This is a difficult operation done only by surgeons in large medical centers. You must be in good overall health with good lung function and no other serious illnesses to tolerate this surgery. Several tests must be done beforehand to be sure you are healthy enough for this surgery. Major complications occur in as many as 1 in 3 people who have this operation.

Pleurectomy/decortication (P/D): This is a less extensive operation in which all of the pleura lining the chest wall (on the side with the cancer) is removed along with the pleura coating the lung on that same side. The pleura coating the mediastinum and the diaphragm is also removed. The lung and diaphragm are not removed.

In a slightly more extensive version of this operation (known as a *radical or extended* P/D), the diaphragm and pericardium are removed as well.

This surgery can be used to try to cure some early cancers, but it is also used as a palliative procedure to relieve symptoms in cases where the entire tumor cannot be removed. It can help control the buildup of fluid, improve breathing, and decrease pain caused by the cancer.

Debulking: The goal of this surgery is to remove as much of the mesothelioma as possible. In general, less tissue is removed in this operation than in a P/D procedure.

Possible side effects of surgery: Possible risks and side effects depend on the extent of the surgery and the person's health beforehand. Serious complications of EPP can include bleeding, blood clots, wound infections, changes in heart rhythm, pneumonia, and loss of lung function. Most of these are less common with less extensive operations.

Because the surgeon must often spread the ribs during surgery, the incision will hurt for some time afterward. Your activity will be limited for at least a month or two.

Surgery for peritoneal mesothelioma

Surgical treatment of peritoneal mesothelioma is often done either to help relieve symptoms or to remove the tumor from the wall of the abdomen and digestive organs. As is the case with pleural mesothelioma, these tumors often have spread too far to be removed completely.

Debulking: The goal of this surgery is to remove as much of the mesothelioma as possible. Sometimes this means removing pieces of the intestine.

After the cancer is debulked (but before the patient is sewn up), chemotherapy may be given into the abdominal cavity. This is called *intraoperative chemotherapy*. If the chemotherapy drugs are heated, it is called *heated intraoperative* (or intraperitoneal) chemotherapy or HIPEC. In either treatment, the drugs are left in a short time, and the patient is sewn up after they are removed.

Omentectomy: The omentum is an apron-like layer of fatty tissue that drapes over the contents inside the abdomen. Cancers in the peritoneum often spread to this tissue, so it may be removed as part of surgery for peritoneal mesothelioma.

Surgery for pericardial mesothelioma

Surgery can remove a mesothelioma from the pericardium (the sac around the heart).

Surgery for mesothelioma of the tunica vaginalis testis

Surgery for mesothelioma of the tunica vaginalis testis, which occurs in the groin, rarely cures this cancer. Most of the time surgery is done because the tumor resembles a hernia. The surgeon attempts to treat a suspected hernia and only realizes the diagnosis after the surgery has begun. This kind of mesothelioma typically can't be removed entirely.

For more information about surgery, please see our document, *Understanding Cancer Surgery: A Guide for Patients and Families*.

Palliative procedures used for malignant mesothelioma

Surgery to remove the cancer is not always an option for patients with malignant mesothelioma. In that case, less invasive procedures can be used to control some of the symptoms caused by mesothelioma, especially those due to built up fluid.

Removal of fluid

Procedures such as thoracentesis, paracentesis, and pericardiocentesis can be used to remove fluid that has built up and is causing symptoms. In these procedures, a doctor uses a long, hollow needle to remove the fluid. These procedures are described in the section, "How is malignant mesothelioma diagnosed?" The major drawback to these techniques is that the fluid often builds up again, so they may need to be repeated.

Pleurodesis

This procedure may be done to try to prevent fluid from building up in the chest. A small cut is made in the skin of the chest wall, and a hollow tube (called a *chest tube*) is placed into the chest so that the fluid can drain out. Then the doctor uses the tube to put a substance into the chest, such as talc mixed in a fluid (talc slurry), the antibiotic doxycycline, or the chemotherapy drug bleomycin. This inflames the linings of the lung (visceral pleura) and chest wall (parietal pleural) so that they stick together, sealing the space and preventing further fluid buildup. The tube is generally left in for a day or two to drain any new fluid that might accumulate. Pleurodesis can also be done at the time of thoracoscopy, either using the drugs doxycycline or bleomycin, or by blowing talc powder into the chest cavity.

Shunt placement

A shunt is a device that allows fluid to move from one part of the body to another. For example, a pleuro-peritoneal shunt lets fluid in the chest move into the abdomen, where it is more likely to be absorbed by the body.

The shunt is a long, thin, flexible tube with a small pump in the middle. In the operating room, the doctor inserts one end of the shunt into the chest cavity and the other end into the peritoneum. (The pump is placed just under the skin over the ribs.) Once the shunt is in place, the patient uses the pump several times a day to move the fluid from the chest to the abdomen. This approach may be used if pleurodesis or other techniques are not effective.

Catheter placement

This is another approach sometimes used to control fluid buildup. One end of the catheter (a thin, flexible tube) is placed in the chest or abdomen through a small cut in the skin, and the other end is left outside the body. This is done in a doctor's office or hospital. Once in place, the catheter can be attached to a special bottle or other device to allow the fluid to drain out on a regular basis.

Radiation therapy for malignant mesothelioma

Radiation therapy uses high-energy x-rays or particles to kill cancer cells. Mesotheliomas are often hard to treat with radiation therapy. They are not usually contained as single, discrete tumors, so aiming radiation at them while avoiding nearby normal tissues is difficult. But new radiation therapy techniques may make this form of treatment more useful.

Uses of radiation therapy

Radiation therapy may be used in different ways to treat mesothelioma:

- It can be used after surgery to try to kill any small areas of cancer that could not be seen and removed during surgery. This is called *adjuvant radiation therapy*.
- It can be used as a palliative procedure to ease symptoms of mesothelioma such as shortness of breath, pain, bleeding, and trouble swallowing.

Types of radiation therapy

Two main types of radiation therapy can be used to treat mesothelioma:

External beam radiation therapy (EBRT): This type of radiation therapy uses x-rays from a machine outside the patient's body to kill cancer cells. It is the most common form of radiation therapy for mesothelioma.

The treatment is much like getting an x-ray, but the radiation is more intense. The procedure itself is painless. Before your treatments start, the medical team will take careful measurements to find the correct angles for aiming the radiation beams and the proper dose of radiation. Each treatment lasts only a few minutes, although the setup time -- getting you into place for treatment -- usually takes longer. Most often, radiation treatments are given 5 days a week for several weeks.

With newer techniques, doctors can treat mesotheliomas more accurately while reducing the radiation reaching nearby healthy tissues such as the lungs. This may offer a better chance of increasing the success rate and reducing side effects.

For example, *intensity-modulated radiation therapy (IMRT)* is an advanced form of 3-dimensional radiation therapy. It uses a computer-driven machine that moves around the patient as it delivers radiation. It shapes the radiation beams to fit the tumor and aims them at the tumor from several angles, as well as adjusting the intensity (strength) of the beams to limit the dose reaching nearby normal tissues.

Brachytherapy: For this type of radiation therapy, a radiation source is placed inside the body, in or near the cancer. The radiation given off travels only a very short distance, which limits the possible damage to nearby healthy tissues. Brachytherapy is seldom used for this type of cancer.

Possible side effects

Side effects of external radiation therapy may include fatigue and sunburn-like skin problems and hair loss where the radiation enters the body. These usually go away once treatment is finished. Chest radiation therapy may cause lung damage and lead to trouble breathing and shortness of breath. Abdominal radiation therapy may cause nausea, vomiting, diarrhea, and a loss of appetite.

If radiation therapy is used together with chemotherapy, the side effects are often worse.

If you are having any side effects from radiation therapy, talk with your doctor. There are often ways to help control these symptoms.

For more general information about radiation therapy, please see our document, *Understanding Radiation Therapy: A Guide for Patients and Families*.

Chemotherapy for malignant mesothelioma

Chemotherapy (chemo) is treatment with anti-cancer drugs. There are 2 main ways that chemotherapy can be given to treat mesothelioma.

In *systemic* therapy, chemotherapy is injected into a vein. The drug enters the bloodstream and travels throughout the body to reach and destroy the cancer cells wherever they may be.

Chemo drugs can also be placed directly into the body cavity where the cancer is – either *intrapleurally* (directly into the chest) or *intraperitoneally* (into the abdomen) – with a small catheter (tube) placed through a small cut in the chest or abdominal wall. Chemo drugs given this way are still absorbed into the bloodstream, but the highest concentration of the drug goes directly to where the cancer cells are.

Chemo drugs are sometimes heated before they are placed directly into a body cavity (called *hyperthermic chemotherapy*), which may help them work better. Sometimes this treatment is given as a single dose in the operating room, right after surgery to remove the cancer. This approach is called *heated intraoperative chemotherapy* or HIPEC. It is more often used to treat peritoneal cancers, in which case it may be called *heated intraperitoneal chemotherapy*.

For mesotheliomas that can be treated with surgery, chemotherapy may be given before surgery to try to shrink the cancer and lower the risk of spread. This is called *neoadjuvant therapy*.

Chemo can also be given after surgery to try to try to kill any cancer cells that were left behind because they were too small to be seen. This type of treatment, called *adjuvant therapy*, may help delay or prevent the cancer growing back.

For cancers that are not resectable, chemotherapy may be the main treatment (alone or along with radiation therapy). Chemotherapy may slow the progression of the disease, but it is very unlikely to make it go away completely.

Doctors usually give chemotherapy in cycles, with each period of treatment followed by a rest period to allow the body time to recover. Chemo cycles generally last about 3 to 4 weeks. Chemotherapy is often not recommended for patients in poor health, but advanced age by itself is not a barrier to getting it.

Several chemo drugs have been used to treat mesothelioma. Most doctors now use a combination of the drugs pemetrexed (Alimta[®]) and cisplatin. Pemetrexed lowers levels of folic acid and vitamin B12 in the body, so patients get these as well to help avoid certain side effects. Other combinations that may be used include pemetrexed with carboplatin and cisplatin with gemcitabine (Gemzar[®]).

Other drugs used to treat mesothelioma include:

- Methotrexate
- Vinorelbine
- Mitomycin
- Doxorubicin (Adriamycin®)
- Epirubicin (Ellence®)
- Cyclophosphamide (Cytoxan®)
- Ifosfamide (Ifex®)

These may be given as combinations of 2 drugs, but single drugs can be used in people who may not be able to tolerate more than one drug. Several other drugs are also being studied for use against mesothelioma.

For HIPEC, either mitomycin or the combination of cisplatin plus doxorubicin are most often used.

Possible side effects

Chemo drugs attack cells that are dividing quickly, which is why they work against cancer cells. But other cells in the body, such as those in the bone marrow, the lining of the mouth and intestines, and the hair follicles, also divide quickly. These cells are also likely to be affected by chemotherapy, which can lead to side effects.

The side effects of chemotherapy depend on the type and dose of drugs you are given and how long they are used for. Common side effects include:

- Hair loss
- Mouth sores
- Loss of appetite
- Nausea and vomiting
- Diarrhea
- Increased chance of infections (from too few white blood cells)
- Easy bruising or bleeding (from too few blood platelets)
- Fatigue (from too few red blood cells)

These side effects are usually short-term and go away after treatment is finished. There are often ways to lessen these side effects. For example, drugs can be given to help prevent or reduce nausea and vomiting. Be sure to ask your doctor or nurse about medicines to help reduce side effects, and let him or her know if you have side effects, so they can be managed effectively.

Some drugs can have other side effects. For example, cisplatin can damage nerves (called *neuropathy*). This can sometimes lead to hearing loss or symptoms in the hands and feet such as pain, burning or tingling sensations, sensitivity to cold or heat, or weakness. In most cases this goes away once treatment is stopped, but it may last a long time in some people.

Be sure to report any side effects or changes you notice while getting chemotherapy to your medical team so that you can get them treated promptly. In some cases, the doses of the drugs may need to be reduced or treatment may need to be delayed or stopped to prevent the effects from getting worse.

For more general information about chemotherapy, please see our document, *Understanding Chemotherapy: A Guide for Patients and Families*. If you'd like more information on a drug used in your treatment or a specific drug mentioned in this section, see our Guide to Cancer Drugs, or call us with the names of the medicines you're taking.

Clinical trials for malignant mesothelioma

You have had to make a lot of decisions since you've been told you have cancer. One of the most important decisions you will make is choosing which treatment is best for you. You may have heard about clinical trials being done for your type of cancer. Or maybe someone on your health care team has mentioned a clinical trial to you.

Clinical trials are carefully controlled research studies that are done with patients who volunteer for them. They are done to get a closer look at promising new treatments or procedures.

If you would like to take part in a clinical trial, you should start by asking your doctor if your clinic or hospital conducts clinical trials. You can also call our clinical trials matching service for a list of clinical trials that meet your medical needs. You can reach this service at 1-800-303-5691 or on our website at www.cancer.org/clinicaltrials. You can also get a list of current clinical trials by calling the National Cancer Institute's Cancer Information Service toll-free at 1-800-4-CANCER (1-800-422-6237) or by visiting the NCI clinical trials website at www.cancer.gov/clinicaltrials.

You must meet requirements to take part in any clinical trial, but if you qualify, it will be up to you whether or not to enter (enroll in) it.

Clinical trials are one way to get state-of-the art cancer treatment. In some cases they may be the only way to get access to newer treatments. Sometimes they may be the only way to get access to some newer treatments. They are also the only way for doctors to learn better methods to treat cancer. Still, they are not right for everyone.

You can get a lot more information on clinical trials in our document called *Clinical Trials: What You Need to Know*. You can read it on our website or call our toll-free number (1-800-227-2345) and have it sent to you.

Complementary and alternative therapies for malignant mesothelioma

When you have cancer you are likely to hear about ways to treat your cancer or relieve symptoms that your doctor hasn't mentioned. Everyone from friends and family to Internet groups and websites may offer ideas for what might help you. These methods can include vitamins, herbs, and special diets, or other methods such as acupuncture or massage, to name a few.

What exactly are complementary and alternative therapies?

Not everyone uses these terms the same way, and they are used to refer to many different methods, so it can be confusing. We use *complementary* to refer to treatments that are used *along with* your regular medical care. *Alternative* treatments are used *instead of* a doctor's medical treatment.

Complementary methods: Most complementary treatment methods are not offered as cures for cancer. Mainly, they are used to help you feel better. Some methods that are used along with regular treatment are: meditation to reduce stress, acupuncture to help relieve pain, or peppermint tea to relieve nausea. Some complementary methods are known to help, while others have not been tested. Some have been proven not be helpful, and a few have even been found harmful.

Alternative treatments: Alternative treatments may be offered as cancer cures. These treatments have not been proven safe and effective in clinical trials. Some of these methods may pose danger, or have life-threatening side effects. But the biggest danger in most cases is that you may lose the chance to be helped by standard medical treatment. Delays or interruptions in your medical treatments may give the cancer more time to grow and make it less likely that treatment will help.

Finding out more

It is easy to see why people with cancer think about alternative methods. You want to do all you can to fight the cancer, and the idea of a treatment with few or no side effects sounds great. Sometimes medical treatments like chemotherapy can be hard to take, or they may no longer be working. But the truth is that most of these alternative methods have not been tested and proven to work in treating cancer.

As you consider your options, here are 3 important steps you can take:

- Look for "red flags" that suggest fraud. Does the method promise to cure all or most cancers? Are you told not to have regular medical treatments? Is the treatment a "secret" that requires you to visit certain providers or travel to another country?
- Talk to your doctor or nurse about any method you are thinking about using.
- Contact us at 1-800-227-2345 to learn more about complementary and alternative methods in general and to find out about the specific methods you are looking at. You

can also learn more on the *Complementary and Alternative Medicine* page of our website.

The choice is yours

Decisions about how to treat or manage your cancer are always yours to make. If you want to use a non-standard treatment, learn all you can about the method and talk to your doctor about it. With good information and the support of your health care team, you may be able to safely use the methods that can help you while avoiding those that could be harmful.

Treatment of mesothelioma based on the extent of the cancer

The stage (extent) of a mesothelioma is an important factor in determining treatment options. But other factors, such as whether the doctor feels the cancer is resectable (all visible cancer can be removed by surgery), as well as a person's general health and preferences, also play a role.

Mesotheliomas can be hard to treat, whether the cancer is resectable or not. It's very important that you understand the goal of treatment before it starts -- whether it is to try to cure the cancer or to help relieve symptoms -- as well as the possible benefits and risks. This can help you make an informed decision when looking at your treatment options.

Resectable mesotheliomas

In general, most stage I and some stage II and III pleural mesotheliomas are potentially resectable, but there are exceptions. Whether a tumor is resectable is also based on the subtype (most doctors don't believe that sarcomatoid tumors are helped by resection), where it is located, how far it has grown into nearby tissues, and whether or not a person is healthy enough to have surgery.

Many patients with resectable pleural mesothelioma have their cancer removed by pleurectomy/decortication (P/D) or extrapleural pneumonectomy (EPP). Surgery is more likely to result in long-term benefit in early stage cancers, where there is a better chance that most or all of the cancer can be removed. For stage I cancers P/D may be done. For other resectable cancers, EPP may be considered if it can be done. It offers the best chance to remove the cancer, but it is a complex and extensive operation and not all patients can tolerate it.

Patients with early-stage peritoneal mesotheliomas might also benefit from surgery that removes as much of the cancer as possible. This may be combined with heated intraoperative chemotherapy (HIPEC). Some patients can have a long remission after this treatment. Surgery may still be helpful for later-stage cancers, but the benefits are more likely to last only a short time.

Sometimes, the surgeon may think the cancer is resectable based on imaging tests (such as CT scans) done before surgery, but once the operation starts it becomes clear that not all of the cancer can be removed. In these cases the surgeon may switch to a less extensive operation like P/D (which is easier to tolerate) or even stop the surgery altogether if it is not likely to be helpful. Treatment would then be the same as for unresectable mesotheliomas (see below).

Doctors are still studying whether giving chemotherapy (chemo) before surgery (neoadjuvant therapy) or giving chemo or radiation therapy after surgery (adjuvant therapy) is helpful. Some doctors prefer to give neoadjuvant chemo, and many doctors advise adjuvant chemo or radiation therapy, but not all doctors agree on what the best course of treatment is.

If you are not healthy enough to have a major operation, you will be treated for unresectable mesothelioma (discussed below).

If you have symptoms because of fluid buildup in the chest or abdomen, other approaches such as thoracentesis/paracentesis or pleurodesis (described in the section about palliative procedures) may be helpful.

Because these cancers can be hard to treat, taking part in a clinical trial of a newer form of treatment may be a reasonable option. These types of studies are usually done in large medical centers.

Unresectable mesotheliomas

Stage IV mesotheliomas, as well as many earlier-stage mesotheliomas, can't be removed completely by surgery. This can be because of the extent or subtype of the cancer or because a person may not be healthy enough for an operation. Chemo is the main treatment for these cancers. It may improve symptoms and shrink or slow the growth of the cancer for a time. Although chemo may help patients live longer, it doesn't cure these cancers. Before starting treatment, the goals of treatment should be clear to you and your family.

In people with early-stage mesotheliomas that are likely to grow slowly and are not causing any symptoms, watching them closely at first may be a reasonable option. Treatment can then be started if there are signs that the cancer is growing quickly or if it starts to cause symptoms.

Because these cancers can be hard to treat, taking part in a clinical trial of a newer form of treatment may be a reasonable option.

In many cases, treatment aimed at relieving symptoms and making you more comfortable may be a good choice. This could include treatments that prevent or reduce fluid buildup in the body, such as thoracentesis/paracentesis or pleurodesis (described in the section about palliative procedures). Sometimes pleurectomy/decortication can help with breathing and pain in the chest.

Pain management is another important aspect of your care. Some minor operations and types of radiation therapy can help relieve pain if needed. Doctors can also prescribe strong pain-relieving drugs. Some people with cancer may hesitate to use opioid drugs (such as morphine) for fear of being sleepy all the time or becoming addicted to them. But many people get very effective pain relief from these medicines without serious side effects. It's very important to let your cancer care team know if you are having pain so that it can be treated effectively.

Recurrent mesotheliomas

Cancer is called *recurrent* when it come backs after treatment. Recurrence can be local (in or near the same place it started) or distant (spread to organs such as the brain or liver). Mesotheliomas often come back after the initial treatment. If this happens, further treatment options depend on where the cancer is, what treatments have already been used, and a person's general health.

In most cases the options will be similar to those listed above for unresectable mesotheliomas. For example, chemo may be used to try to shrink or slow the growth of the cancer and to relieve any symptoms. Because recurrent cancers can often be hard to treat, clinical trials of new types of treatment may be a good option. For more information on how to deal with a recurrence, you may want to read our document, *When Your Cancer Comes Back: Cancer Recurrence*.

More treatment information for malignant mesothelioma

For more details on treatment options -- including some that may not be addressed in this document -- the National Cancer Institute (NCI) and the National Comprehensive Cancer Network (NCCN) are good sources of information.

The NCI provides treatment guidelines via its telephone information center (1-800-4-CANCER) and its website (www.cancer.gov). Detailed guidelines intended for use by cancer care professionals are also available on www.cancer.gov.

The NCCN, made up of experts from many of the nation's leading cancer centers, develops cancer treatment guidelines for doctors to use when treating patients. These are available on the NCCN website (www.nccn.org).

What should you ask your doctor about malignant mesothelioma?

As you cope with cancer and its treatment, you need to have honest, open discussions with your doctor. You should feel free to ask any question, no matter how small it might seem. Here are some questions you might want to ask. Nurses, social workers, and other members of the treatment team may also be able to answer many of your questions.

• What kind of mesothelioma do I have?

- Has my cancer spread beyond where it started?
- What is my cancer's stage (extent), and what does that mean?
- Is my cancer likely to be resectable (removable by surgery)?
- Will I need other tests before we can decide on treatment?
- Will I need to see other doctors?
- How much experience do you have treating this type of cancer?
- Should I get a second opinion?
- What are my treatment options?
- What is the goal of treatment?
- What do you recommend and why?
- What risks or side effects are there to the treatments you suggest?
- What should I do to be ready for treatment?
- How long will treatment last? What will it be like? Where will it be done?
- How will treatment affect my daily activities?
- What will we do if the treatment doesn't work or if the cancer recurs?
- What type of follow-up might I need after treatment?

Along with these sample questions, be sure to write down some of your own. For instance, you might want more information about recovery times. Or you may want to ask if you qualify for any clinical trials.

What happens after treatment for malignant mesothelioma?

For some people with mesothelioma, treatment may remove or destroy the cancer. Completing treatment can be both stressful and exciting. You may be relieved to finish treatment, but find it hard not to worry about cancer growing or coming back. (When cancer comes back after treatment, it is called *recurrence*.) This is a very common concern in people who have had cancer.

It may take a while before your fears lessen. But it may help to know that many cancer survivors have learned to live with this uncertainty and are leading full lives. Our document, *Living With Uncertainty: The Fear of Cancer Recurrence*, gives more detailed information on this.

For many people, the mesothelioma may never go away completely. These people may get regular treatments with chemotherapy, radiation therapy, or other therapies to help keep the cancer in check. Learning to live with cancer that doesn't go away can be difficult and very stressful. It has its own type of uncertainty. Our document *When Cancer Doesn't Go Away* talks more about this.

Follow-up care

If you have completed treatment, your doctors will still want to watch you closely. It is very important to keep all follow-up appointments. During these visits, your doctors will ask about symptoms, do physical exams, and may order blood tests (such as the osteopontin or MesoMark tests) or imaging tests such as CT scans or PET scans. There is no widely agreed upon follow-up schedule for people with mesothelioma. Your doctor will most likely want to see you fairly often (at least every 3 months or so) at first. The time between visits may be extended if there are no problems.

Follow-up is needed to check for cancer recurrence or spread, as well as possible side effects of certain treatments. This is the time for you to ask your health care team any questions you might have and to discuss any concerns.

Almost any cancer treatment can have side effects. Some may last for a few weeks to several months, but others can be permanent. Don't hesitate to tell your cancer care team about any symptoms or side effects that bother you so they can help you manage them.

If the cancer does recur, further treatment will depend on the location of the cancer, what treatments you've had before, and your health. For more information on how recurrent cancer is treated, see the section "Treatment of mesothelioma based on the extent of the cancer." For more general information on dealing with a recurrence, you may also want to read our document, *When Your Cancer Comes Back: Cancer Recurrence*. You can get this document by calling 1-800-227-2345.

Seeing a new doctor

At some point after your cancer diagnosis and treatment, you may find yourself seeing a new doctor who does not know anything about your medical history. It is important that you be able to give your new doctor the details of your diagnosis and treatment. Gathering these details soon after treatment may be easier than trying to get them at some point in the future. Make sure you have the following information handy:

- A copy of your pathology report(s) from any biopsies or surgeries
- If you had surgery, a copy of your operative report(s)
- If you stayed in the hospital, a copy of the discharge summary that doctors prepare when patients are sent home
- If you had radiation therapy, a summary of the type and dose of radiation and when and where it was given

• If you had chemotherapy, a list of your drugs, drug doses, and when you took them

It is also important to keep health insurance. Tests and doctor visits cost a lot, and even though no one wants to think of their cancer coming back, this could happen. For more information about health insurance, see our document called *Health Insurance and Financial Assistance for the Person With Cancer*.

Lifestyle changes after malignant mesothelioma

You can't change the fact that you have had cancer. What you can change is how you live the rest of your life -- making choices to help you stay healthy and feel as well as you can. This can be a time to look at your life in new ways. Maybe you are thinking about how to improve your health over the long term. Some people even start during cancer treatment.

Make healthier choices

For many people, a diagnosis of cancer helps them focus on their health in ways they might not have thought much about in the past. Are there things you could do that might make you healthier? Maybe you could try to eat better or get more exercise. Maybe you could cut down on the alcohol, or give up tobacco. Even things like keeping your stress level under control may help. Now is a good time to think about making changes that can have positive effects for the rest of your life. You will feel better and you will also be healthier.

You can start by working on those things that worry you most. Get help with those that are harder for you. For instance, if you are thinking about quitting smoking and need help, call the American Cancer Society at 1-800-227-2345.

Eating better

Eating right can be hard for anyone, but it can get even tougher during and after cancer treatment. Treatment may change your sense of taste. Nausea can be a problem. You may not feel like eating and lose weight when you don't want to. Or you may have gained weight that you can't seem to lose. All of these things can be very frustrating.

If treatment caused weight changes or eating or taste problems, do the best you can and keep in mind that these problems usually get better over time. You may find it helps to eat small portions every 2 to 3 hours until you feel better. You may also want to ask your cancer team about seeing a dietitian, an expert in nutrition who can give you ideas on how to deal with these treatment side effects.

One of the best things you can do after cancer treatment is put healthy eating habits into place. You may be surprised at the long-term benefits of some simple changes, like increasing the variety of healthy foods you eat. Getting to and staying at a healthy weight, eating a healthy diet, and limiting your alcohol intake may lower your risk for a number of types of cancer, as well as having many other health benefits. You can read more about

this in our document, Nutrition and Physical Activity During and After Cancer Treatment: Answers to Common Questions.

Rest, fatigue, and exercise

Extreme tiredness, called *fatigue*, is very common in people treated for cancer. This is not a normal tiredness, but a bone-weary exhaustion that doesn't get better with rest. For some people, fatigue lasts a long time after treatment, and can make it hard for them to exercise and do other things they want to do. But exercise can help reduce fatigue. Studies have shown that patients who follow an exercise program tailored to their personal needs feel better physically and emotionally and can cope better, too.

If you were sick and not very active during treatment, it is normal for your fitness, endurance, and muscle strength to decline. Any plan for physical activity should fit your own situation. A person who has never exercised will not be able to take on the same amount of exercise as someone who plays tennis twice a week. If you haven't exercised in a few years, you will have to start slowly -- maybe just by taking short walks.

Talk with your health care team before starting anything. Get their opinion about your exercise plans. Then, try to find an exercise buddy so you're not doing it alone. Having family or friends involved when starting a new exercise program can give you that extra boost of support to keep you going when the push just isn't there.

If you are very tired, you will need to balance activity with rest. It is OK to rest when you need to. Sometimes it's really hard for people to allow themselves to rest when they are used to working all day or taking care of a household, but this is not the time to push yourself too hard. Listen to your body and rest when you need to. (For more information on dealing with fatigue, please see *Fatigue in People With Cancer* and *Anemia in People With Cancer*.)

Keep in mind exercise can improve your physical and emotional health.

- It improves your cardiovascular (heart and circulation) fitness.
- Along with a good diet, it will help you get to and stay at a healthy weight.
- It makes your muscles stronger.
- It reduces fatigue and helps you have more energy.
- It can help lower anxiety and depression.
- It can make you feel happier.
- It helps you feel better about yourself.

And long term, we know that getting regular physical activity plays a role in helping to lower the risk of some cancers, as well as having other health benefits.

How does having malignant mesothelioma affect your emotional health?

During and after treatment, you may find yourself overcome with many different emotions. This happens to a lot of people.

You may find yourself thinking about death and dying. Or maybe you're more aware of the effect the cancer has on your family, friends, and career. You may take a new look at your relationships with those around you. Unexpected issues may also cause concern. For instance, you may see your health care team less often after treatment and have more time on your hands. These changes can make some people anxious.

Almost everyone who is going through or has been through cancer can benefit from getting some type of support. You need people you can turn to for strength and comfort. Support can come in many forms: family, friends, cancer support groups, church or spiritual groups, online support communities, or one-on-one counselors. What's best for you depends on your situation and personality. Some people feel safe in peer-support groups or education groups. Others would rather talk in an informal setting, such as church. Others may feel more at ease talking one-on-one with a trusted friend or counselor. Whatever your source of strength or comfort, make sure you have a place to go with your concerns.

The cancer journey can feel very lonely. It is not necessary or good for you to try to deal with everything on your own. And your friends and family may feel shut out if you do not include them. Let them in, and let in anyone else who you feel may help. If you aren't sure who can help, call your American Cancer Society at 1-800-227-2345 and we can put you in touch with a group or resource that may work for you. You may also want to read our booklet *Distress in People with Cancer*, which is available online, or you can call us to request a free copy by mail.

If treatment of malignant mesothelioma stops working

If cancer keeps growing or comes back after one kind of treatment, it may be possible to try another treatment plan that might still cure the cancer, or at least shrink the tumors enough to help you live longer and feel better. But when many different treatments have been tried and the cancer has not gotten any better, the cancer tends to become resistant to all treatment. If this happens, it's important to weigh the possible limited benefits of a new treatment against the possible downsides, including treatment side effects. Everyone has their own way of looking at this.

This is likely to be the hardest part of your battle with cancer -- when you have been through many medical treatments and nothing's working anymore. Your doctor might be able to offer you new options, but at some point you need to consider that treatment is not likely to improve your health or change your outcome or survival.

If you want to continue to get treatment for as long as you can, you need to think about the odds of treatment having any benefit and how this compares to the possible risks and side effects. In many cases, your doctor can estimate how likely it is the cancer will respond to treatment you are considering. For instance, the doctor may say that more treatment might have about a 1 in 100 chance of working. Some people are still tempted to try this. But it is important to think about and understand your reasons for choosing this plan.

No matter what you decide to do, it is important that you feel as good as you can. Make sure you are asking for and getting treatment for any symptoms you might have, such as nausea or pain. This type of treatment is called *palliative care*.

Palliative care helps relieve symptoms, but it is not expected to cure the disease. It can be given along with cancer treatment, or can even be cancer treatment. The difference is its purpose -- the main goal of palliative care is to improve the quality of your life, or help you feel as good as you can for as long as you can. Sometimes this means using drugs to help with symptoms like pain or nausea. Sometimes, though, the treatments used to control your symptoms are the same as those used to treat cancer. For instance, radiation might be used to help relieve bone pain caused by cancer that has spread to the bones. Or chemo might be used to help shrink a tumor and keep it from blocking the bowels. But this is not the same as treatment to try to cure the cancer. You can learn more about the changes that occur when curative treatment stops working, and about planning ahead for yourself and your family, in our documents called *Nearing the End of Life* and *Advance Directives*. You can read them online or call us to have free copies mailed to you.

At some point, you may benefit from hospice care. This is special care that treats the person rather than the disease; it focuses on quality rather than length of life. Most of the time, it is given at home. Your cancer may be causing problems that need to be managed, and hospice focuses on your comfort. You should know that while getting hospice care often means the end of treatments such as chemo and radiation, it doesn't mean you can't have treatment for the problems caused by your cancer or other health conditions. In hospice the focus of your care is on living life as fully as possible and feeling as well as you can at this difficult time. You can learn more about hospice in our document called *Hospice Care*.

Staying hopeful is important, too. Your hope for a cure may not be as bright, but there is still hope for good times with family and friends – times that are filled with happiness and meaning. Pausing at this time in your cancer treatment gives you a chance to refocus on the most important things in your life. Now is the time to do some things you've always wanted to do and to stop doing the things you no longer want to do. Though the cancer may be beyond your control, there are still choices you can make.

What's new in malignant mesothelioma research and treatment?

There is always research going on in the area of mesothelioma. Scientists are looking for ways to prevent, diagnose, and treat mesothelioma. Despite recent progress, much remains to be learned about the best way to treat these cancers.

Causes and prevention

Much of the research on mesothelioma has focused on learning exactly how asbestos changes mesothelial cells and their DNA to cause these cancers. Understanding how these fibers produce cancer might help us develop ways to prevent those changes.

The role of asbestos in increasing the risk of mesothelioma is a definite public health concern. Researchers are learning more about which fibers can produce cancer, how they cause these cancers, and what levels of exposure can be considered safe. Now that the dangers of asbestos are known, we can limit or stop exposure in homes, public buildings, and the workplace. Unfortunately, regulations protecting workers from asbestos exposure are much less stringent in some countries than in others.

Research is also under way to clarify the role (if any) of SV40, a virus that has been linked to mesothelioma in some studies.

Treatment

Mesothelioma remains a difficult cancer to treat, and doctors are constantly trying to improve on current approaches. The exact roles of surgery, radiation therapy, and chemotherapy in the treatment of mesothelioma are highly debated. Combinations of these treatments, called *multimodality therapy*, are now being tested and may provide the most promising option for some patients. Newer types of treatment now being studied may give patients and their doctors even more options.

Chemotherapy

Some chemotherapy drugs can shrink or slow the growth of mesotheliomas, but in most cases the effects last for a limited time. Studies are underway to test newer chemotherapy drugs.

Photodynamic therapy

Another technique now being studied is photodynamic therapy (PDT). For this treatment, a light-activated drug is injected into a vein. The drug spreads throughout the body and tends to collect in cancer cells. A few days later (usually just after surgery for the mesothelioma), a special red light on the end of a tube is placed into the chest cavity. The light causes a chemical change that activates the drug and causes the cancer cells to die. Since the drug is only active in the areas exposed to the special light, this approach may cause fewer side effects than use of drugs that spread throughout all tissues of the body. Several clinical trials are now studying the use of PDT for mesothelioma. You can find out more about PDT in our document, *Photodynamic Therapy*.

Targeted drugs

In general, chemotherapy drugs are limited in their effectiveness against mesothelioma. As researchers have learned more about the changes in cells that cause cancer, they have been able to develop newer drugs that specifically target these changes. Targeted drugs work differently from standard chemotherapy drugs. They may sometimes work when chemotherapy drugs do not, and they often have different (and less severe) side effects.

Sunitinib (Sutent) is an example of a targeted drug that has shown promise in studies. .

Other new drugs have different targets. For example, some new drugs target mesothelin, a protein found in high levels in mesothelioma cells. You can learn more about targeted therapy drugs in our document, *Targeted Therapy*.

Newer forms of treatment

Because standard treatments often have limited usefulness against mesothelioma, researchers are studying other new types of treatment as well.

Gene therapy: A newer type of treatment being tested on mesothelioma is *gene therapy*, which attempts to add new genes to cancer cells to make them easier to kill. One approach to gene therapy uses special viruses that have been modified in the lab. The virus is injected into the pleural space and infects the mesothelioma cells. When this infection occurs, the virus injects the desired gene into the cells. In one version of this approach, the virus carries a gene that helps turn on the immune system to attack the cancer cells. Early studies of this approach have found it may shrink or slow the growth of mesothelioma in some people, but more research is needed to confirm this.

Immunotherapy: Other new treatments called *cancer vaccines* are also aimed at getting the immune system to attack the cancer. In one approach, immune cells are removed from a patient's blood and treated in the lab to get them to react to tumor cells. The immune cells are then given back to the patient as blood transfusions, where it is hoped they will cause the body's immune system to attack the cancer. This approach is now being studied in clinical trials.

Another form of immunotherapy that is being studied is a drug called tremelimumab that targets a certain white blood cell and takes the brakes off the immune system.

You can learn more about this topic in our document *Immunotherapy*.

Additional resources for malignant mesothelioma

More information from your American Cancer Society

Here is more information you might find helpful. You also can order free copies of our documents from our toll-free number, 1-800-227-2345, or read them on our website, www.cancer.org.

Living with cancer

After Diagnosis: A Guide for Patients and Families (also available in Spanish)

Living With Uncertainty: The Fear of Cancer Recurrence

Clinical Trials: What You Need to Know (also available in Spanish)

Pain Control: A Guide for Those With Cancer and Their Loved Ones (also available in

Spanish)

Nutrition and Physical Activity During and After Cancer Treatment: Answers to

Common Questions

When Your Cancer Comes Back: Cancer Recurrence

Understanding cancer treatments

Understanding Cancer Surgery: A Guide for Patients and Their Families (also available in Spanish)

Understanding Radiation Therapy: A Guide for Patients and Their Families (also available in Spanish)

Understanding Chemotherapy: A Guide for Patients and Their Families (also available in Spanish)

Family and caregiver concerns

Talking With Friends and Relatives About Your Cancer (also in Spanish)

What It Takes to Be a Caregiver

Caring for the Patient With Cancer at Home: A Guide for Patients and Families (also available in Spanish)

Helping Children When a Family Member Has Cancer: Dealing With Diagnosis (also available in Spanish)

Work, insurance, and finances

Health Insurance and Financial Assistance for the Cancer Patient

Returning to Work After Cancer Treatment

Working During Cancer Treatment

Carcinogens and cancer

Asbestos (also available in Spanish)

Questions About Smoking, Tobacco, and Health (also available in Spanish)

Guide to Quitting Smoking (also available in Spanish)

Testing and diagnosis of malignant mesothelioma

Endoscopy

Testing Biopsy and Cytology Specimens for Cancer

When treatment stops working

Nearing the End of Life

Advance Directives

Hospice Care

Your American Cancer Society also has books that you might find helpful. Call us at 1-800-227-2345 or visit our bookstore online at cancer.org/bookstore to find out about costs or to place an order.

National organizations and websites*

In addition to the American Cancer Society, other sources of patient information and support include:

Information on asbestos

Agency for Toxic Substances and Disease Registry

Toll-free number: 1-800-232-4636 Website: www.atsdr.cdc.gov/

Offers information on toxic substances, including a special section on asbestos with fact sheets and video information

Environmental Protection Agency

Telephone: 1-202-272-0167 Website: www.epa.gov/

Information about asbestos at http://www2.epa.gov/asbestos

Occupational Safety and Health Administration

Toll-free number: 1-800-321-6742 (1-800-321-OSHA)

Website: www.osha.gov/

Addresses safety and health at work; contains a fact sheet on asbestos at https://www.osha.gov/OshDoc/data_AsbestosFacts/asbestos-factsheet.pdf; also has an entire section on asbestos at https://www.osha.gov/SLTC/asbestos/

Information on mesothelioma and cancer in general

Mesothelioma Applied Research Foundation

Toll-free number: 1-877-363-6376 (1-877-END-MESO)

Website: www.curemeso.org

More on mesothelioma, peer support via other patients and families dealing with mesothelioma, and referrals to clinical trials and doctors who specialize in mesothelioma. Booklet: "100 Questions and Answers About Mesothelioma"

National Cancer Institute

Telephone: 1-800-422-6237 (1-800-4-CANCER)

Website: www.cancer.gov

Provides information on all types of cancer, living with cancer, support information for families of people with cancer, research, and more

National Coalition for Cancer Survivorship

Toll-free number: 1-888-650-9127

1-877-NCCS-YES (622-7937) for some publications and Cancer Survivor Toolbox®

orders

Website: www.canceradvocacy.org

Offers information on work, health insurance, and more. The Cancer Survival Toolbox is a free, self-learning audio program to help cancer survivors and caregivers develop practical tools needed to deal with the diagnosis, treatment and challenges of cancer. Listen online or order CDs. Also in Spanish and Chinese

No matter who you are, we can help. Contact us anytime, day or night, for information and support. Call us at **1-800-227-2345** or visit www.cancer.org.

References: Malignant mesothelioma detailed guide

Alexander HR Jr, Bartlett DL, Pingpank JF, et al. Treatment factors associated with long-term survival after cytoreductive surgery and regional chemotherapy for patients with malignant peritoneal mesothelioma. *Surgery*. 2013 Jun;153(6):779-786. Epub 2013 Mar 13.

American Joint Committee on Cancer. Pleural mesothelioma. *AJCC Cancer Staging Manual*. 7th ed. New York, NY: Springer; 2010: 271-274.

Arrieta Ó, Medina LA, Estrada-Lobato E, Hernández-Pedro N, Villanueva-Rodríguez G, Martínez-Barrera L, Macedo EO, López-Rodríguez V, Motola-Kuba D, Corona-Cruz JF. First-line chemotherapy with liposomal doxorubicin plus cisplatin for patients with

^{*}Inclusion on this list does not imply endorsement by the American Cancer Society.

advanced malignant pleural mesothelioma: phase II trial. *Br J Cancer*. 2012 Mar 13;106(6):1027-1032. Epub 2012 Feb 21.

Calabrò L, Morra A, Fonsatti E, et al. Tremelimumab for patients with chemotherapyresistant advanced malignant mesothelioma: an open-label, single-arm, phase 2 trial. *Lancet Oncol*. 2013 Oct;14(11):1104-1111. Epub 2013 Sep 11.

Howlader N, Noone AM, Krapcho M, et al (eds). SEER Cancer Statistics Review, 1975-2010, National Cancer Institute. Bethesda, MD, http://seer.cancer.gov/csr/1975_2010/, based on November 2012 SEER data submission, posted to the SEER web site, April 2013.

Hegmans JP, Veltman JD, Lambers ME, et al. Consolidative dendritic cell-based immunotherapy elicits cytotoxicity against malignant mesothelioma. *Am J Respir Crit Care Med*. 2010;181:1383-1390.

Mirarabshahii P, Pillai K, Chua TC, Pourgholami MH, Morris DL. Diffuse malignant peritoneal mesothelioma--an update on treatment. *Cancer Treat Rev.* 2012 Oct;38(6):605-612. Epub 2011 Nov 21.

National Cancer Institute. Physician Data Query (PDQ). Malignant Mesothelioma: Treatment. 7/16/2012. Accessed at www.cancer.gov/cancertopics/pdq/treatment/malignantmesothelioma/HealthProfessional on September 5, 2013.

National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Malignant Pleural Mesothelioma. V.2.1014. Accessed at www.nccn.org/professionals/physician_gls/pdf/mpm.pdf on September 5, 2013.

Nowak AK, Millward MJ, Creaney J, Francis RJ, Dick IM, Hasani A, van der Schaaf A, Segal A, Musk AW, Byrne MJ. A phase II study of intermittent sunitinib malate as second-line therapy in progressive malignant pleural mesothelioma. *J Thorac Oncol*. 2012 Sep;7(9):1449-1456.

Pan X, Day W, Wang W, et al. Residential proximity to naturally occurring asbestos and mesothelioma risk in California. *Am J Resp Crit Care*. 2005;172:1019-1025.

Pass HI, Vogelzgang NJ, Hahn SM, Carbone M. Benign and malignant mesothelioma. In: DeVita VT, Lawrence TS, Rosenberg SA, eds. *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology.* 9th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2011:2052-2080.

Price B, Ware A. Mesothelioma trends in the United States: An update based on Surveillance Epidemiology and End Results program date for 1973-2003. *Am J Epidemiol*. 2004;159:107-112.

Robinson BWS, Lake RA. Advances in malignant mesothelioma. *N Engl J Med*. 2005;353:1591-1603.

Rusch VW, Giroux D, Kennedy C, Ruffini E, Cangir AK, Rice D, Pass H, Asamura H, Waller D, Edwards J, Weder W, Hoffmann H, van Meerbeeck JP; IASLC Staging

Committee. Initial analysis of the international association for the study of lung cancer mesothelioma database. *J Thorac Oncol*. 2012 Nov;7(11):1631-1639.

Sovak MA, Aisner SC, Aisner J. Tumors of the pleura and mediastinum. In: Abeloff MD, Armitage JO, Niederhuber JE. Kastan MB, McKenna WG, eds. *Abeloff's Clinical Oncology*. 4th ed. Philadelphia, Pa: Elsevier; 2008:1367-1398.

Sterman DH, Haas A, Moon E, et al. A trial of intrapleural adenoviral-mediated interferon-α2b gene transfer for malignant pleural mesothelioma. *Am J Respir Crit Care Med.* 2011;184:1395-1399.

Testa JR, Cheung M, Pei J, et al. Germline BAP1 mutations predispose to malignant mesothelioma. *Nat Genet*. 2011;43:1022-1025.

Tsao AS, Wistuba I, Roth JA, Kindler HL. Malignant pleural mesothelioma. *J Clin Oncol*. 2009;27:2081-2090.

Last Medical Review: 12/19/2013

Last Revised: 12/19/2013

2013 Copyright American Cancer Society

For additional assistance please contact your American Cancer Society 1-800-227-2345 or www.cancer.org