



Breast Cancer Fact Sheet

for Health Care Professionals

Breast Cancer in the US^{1, 2}

In women, breast cancer is the most common cancer diagnosed (after skin cancer) and the second-leading cause of cancer death (after lung cancer).

Types of Breast Cancer^{1, 3}

There are several types of breast cancer, including: ductal carcinoma in situ (DCIS), invasive ductal carcinoma (IDC), invasive lobular carcinoma (ILC), triple-negative breast cancer (TNBC), inflammatory breast cancer (IBC), angiosarcoma of the breast, Phyllodes tumor, and Paget disease of the breast.

Risk Factors^{1, 2, 3}

Sex: Both men and women can develop this disease, but being born female is the main risk factor for breast cancer.

Age: The risk of developing breast cancer increases with age, and most breast cancers are found in women ages 55 or older.

Personal or family history: Breast cancer risk is higher among women with a personal or family history of the disease.

- About 5 to 10% of breast cancers are likely caused by genetic mutations, such as those in the *BRCA1* and *BRCA2* genes.
- Having a first-degree relative with breast cancer increases risk, while having more than one first-degree relative who has or had breast cancer increases the risk even more. Risk is further increased when the affected female relative was diagnosed at a young age or was diagnosed with cancer in both breasts, or if the affected relative is male.
- Previous history of breast cancer or certain benign breast conditions, such as atypical hyperplasia, can increase risk.

Other Risk Factors^{1, 2, 3}

- Drinking alcohol
- Physical inactivity
- Having dense breast tissue



- Using postmenopausal hormone therapy with estrogen and progesterone therapy
- Weight gain after age 18 or having excess body weight (for postmenopausal breast cancer)
- A long menstrual history – starting menstruation early or having late menopause
- Not having children, not breastfeeding, or having first live birth after age 30
- Using certain birth control methods
- Undergoing radiation to the chest before age 30

Screening and Detection^{1, 4, 5}

The American Cancer Society recommends the following guidelines for the early detection of breast cancer in average-risk women:

- **Women ages 40 to 44** should have the choice to start annual breast cancer screening with mammograms.
- **Women ages 45 to 54** should get mammograms every year.
- **Women 55 and older** can switch to mammograms every 2 years, or can continue yearly screening.
- Screening should continue as long as a woman is in good health and is expected to live at least 10 more years.
- **All women** should be familiar with the known benefits, limitations, and potential harms linked to breast cancer screening.
- A screening MRI is recommended for **women at high risk of breast cancer**, including women with a strong family history of breast or ovarian cancer; those with a lifetime risk of breast cancer of about 20% to 25% or greater, according to risk assessment tools that are based mainly on family history; those with a known breast cancer gene mutation; and women who were treated with radiation therapy to the chest before age 30.

Signs and Symptoms^{1, 3}

The most common physical sign of breast cancer is a new, painless lump or mass. Sometimes breast cancer spreads to axillary lymph nodes and causes a lump or swelling, even before the original breast tumor is large enough to be felt. Less common signs and symptoms include breast pain or heaviness; persistent changes, such as swelling, thickening, redness, irritation, or dimpling of the skin; and nipple changes, such as spontaneous discharge (especially if bloody), pain, retraction, redness, or scaliness. Any persistent change in the breast should be evaluated by a physician.

Prevention^{1, 4}

There is no sure way to prevent breast cancer, and some risk factors can't be changed, such as age, race, family history of disease, genetic mutations, and reproductive history. Lifestyle factors, such as avoiding or limiting alcohol, breastfeeding, engaging in regular physical activity, and staying at a healthy weight, are associated with lower risk. Selective estrogen receptor modulators (SERMs), such as tamoxifen and raloxifene, and prophylactic mastectomy can help reduce the risk in some high-risk women.

Treatment^{1, 3, 5}

Treatment options are based on the tumor subtype, stage, and molecular characteristics, along with patient comorbidities. Surgery, radiation therapy, chemotherapy, targeted therapy, and immunotherapy drugs, either in combination or alone, are common treatments that might be used. Visit [cancer.org/cancer/breast-cancer](https://www.cancer.org/cancer/breast-cancer) to learn more about treatment options for the different types of breast cancer.

References

1. American Cancer Society. *Breast Cancer Facts & Figures 2024-2025*. Atlanta: American Cancer Society; 2024. Accessed at <https://www.cancer.org/research/cancer-facts-statistics/breast-cancer-facts-figures.html> on January 22, 2025.
2. American Cancer Society. *Cancer Facts & Figures 2025*. Atlanta: American Cancer Society; 2025. Accessed at <https://www.cancer.org/research/cancer-facts-statistics/all-cancer-facts-figures/2025-cancer-facts-figures.html> on January 22, 2025.
3. American Cancer Society. *Breast Cancer 2025*. Accessed at <https://www.cancer.org/cancer/breast-cancer.html> on January 22, 2025.
4. Smith RA, Andrews KS, Brooks D, et. al. Cancer screening in the United States, 2019: A review of current American Cancer Society guidelines and current issues in cancer screening. *CA Cancer J Clin*. 2019; 69:184-210.
5. National Comprehensive Cancer Network. Clinical Practice Guidelines in Oncology (NCCN Guidelines). Breast Cancer, Version 6.2024. Accessed at https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf on January 22, 2025.
6. American Cancer Society. *Cancer Treatment & Survivorship Facts & Figures, 2022-2024*. Atlanta: American Cancer Society. Accessed at <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/cancer-treatment-and-survivorship-facts-and-figures/2022-cancer-treatment-and-survivorship-fandf-acf.pdf> on January 22, 2025.
7. Edwards RL, Taylor RA, Bakitas MA. Integration of palliative care. In: Haylock PJ, Curtiss CP, eds. *Cancer Survivorship: Interprofessional, Patient-Centered Approaches to the Seasons of Survival*. Oncology Nursing Society; 2019: 137-151.

Breast Cancer in the US:

2025 estimates^{1, 2}

New cases:

- Invasive breast cancer in women: 316,950
- Ductal carcinoma in situ (DCIS) in women: 59,080
- Invasive breast cancer in men: 2,800

Deaths: 42,680

- Women: 42,170
- Men: 510

5-year relative survival rate for localized stages: 99%

5-year relative survival rate for all stages combined: 91%

Quality of Life^{2, 3, 6, 7}

Common issues affecting quality of life for patient with breast cancer include uncertainty about treatment options, changes in physical appearance (including hair loss and post surgical effects), concerns about side effects (such as lymphedema, hot flashes, pain, fatigue, depression, sleep difficulties, and sexual and fertility changes), guilt for things that may have contributed to cancer diagnosis, fear of recurrence, changes in lifestyle following treatment, and the burden on finances and loved ones.

A cancer diagnosis can profoundly impact quality of life. **Clinicians should assess for any physical, social, psychological, spiritual, and financial issues.** Integrating palliative care can help manage symptoms, address issues, and improve quality of life. It can be offered at any time, from the point of diagnosis through treatment, and until the end of life. Throughout a person's cancer journey, it's very important for clinicians to share information and coordinate care to ensure surveillance is ongoing.

