

Estimated 1-5 Year Cancer Survivors 2012

STATE	2012	1 Year	2011	2 Year	2010	3 Year	2009	4 Year	2008	5 Year	2007	Estimated
	Estimated	Survival	Estimated	Survival	Estimated	Survival	Estimated	Survival	Estimated	Survival	Estimated	1-5 Year
	Cases		Cases		Cases		Cases		Cases		Cases	Survivors
AL	26,440	77.1%	25,530	68.9%	23,640	63.9%	24,090	60.1%	22,340	57.0%	20,590	76,530
AK	3,640	77.1%	3,090	68.9%	2,860	63.9%	2,530	60.1%	2,650	57.0%	2,500	8,990
AZ	31,990	77.1%	31,550	68.9%	29,780	63.9%	27,600	60.1%	25,540	57.0%	26,270	92,800
AR	16,120	77.1%	16,070	68.9%	15,320	63.9%	14,800	60.1%	14,840	57.0%	14,130	49,380
CA	165,810	77.1%	163,480	68.9%	157,320	63.9%	152,170	60.1%	156,530	57.0%	151,250	511,960
CO	22,820	77.1%	22,390	68.9%	21,340	63.9%	20,340	60.1%	18,900	57.0%	19,190	67,260
CT	21,530	77.1%	21,440	68.9%	20,750	63.9%	20,650	60.1%	19,190	57.0%	19,780	66,830
DE	5,340	77.1%	5,130	68.9%	4,890	63.9%	4,690	60.1%	4,590	57.0%	4,530	15,660
DC	2,980	77.1%	2,830	68.9%	2,760	63.9%	2,600	60.1%	2,560	57.0%	2,540	8,730
FL	117,580	77.1%	113,400	68.9%	107,000	63.9%	102,210	60.1%	101,920	57.0%	106,560	348,460
GA	48,130	77.1%	44,580	68.9%	40,480	63.9%	39,080	60.1%	36,980	57.0%	35,440	129,660
HI	6,610	77.1%	6,710	68.9%	6,670	63.9%	6,400	60.1%	6,310	57.0%	6,020	21,080
ID	7,720	77.1%	7,520	68.9%	7,220	63.9%	6,800	60.1%	6,430	57.0%	6,140	22,480
IL	65,750	77.1%	65,610	68.9%	63,890	63.9%	60,960	60.1%	59,130	57.0%	62,010	204,440
IN	35,060	77.1%	34,050	68.9%	33,020	63.9%	31,320	60.1%	29,550	57.0%	30,040	103,900
IA	17,010	77.1%	17,500	68.9%	17,260	63.9%	16,740	60.1%	16,150	57.0%	16,540	55,220
KS	14,090	77.1%	14,070	68.9%	13,550	63.9%	13,080	60.1%	12,520	57.0%	12,760	43,340
KY	25,160	77.1%	25,010	68.9%	24,240	63.9%	24,060	60.1%	23,270	57.0%	22,850	78,370
LA	23,480	77.1%	22,780	68.9%	20,950	63.9%	22,170	60.1%	23,360	57.0%	22,540	73,050
ME	8,990	77.1%	8,820	68.9%	8,650	63.9%	9,000	60.1%	8,140	57.0%	8,340	28,160
MD	31,000	77.1%	28,890	68.9%	27,700	63.9%	26,650	60.1%	27,380	57.0%	26,390	89,890
MA	38,470	77.1%	37,470	68.9%	36,040	63.9%	36,080	60.1%	34,470	57.0%	34,920	117,400
MI	57,790	77.1%	57,010	68.9%	55,660	63.9%	53,550	60.1%	51,150	57.0%	54,410	178,280
MN	28,060	77.1%	27,600	68.9%	25,080	63.9%	23,670	60.1%	23,160	57.0%	25,420	82,090
MS	15,190	77.1%	14,990	68.9%	14,330	63.9%	14,150	60.1%	13,400	57.0%	12,470	45,630
MO	33,440	77.1%	32,740	68.9%	31,160	63.9%	30,090	60.1%	29,390	57.0%	29,930	100,660
MT	5,550	77.1%	5,690	68.9%	5,570	63.9%	5,340	60.1%	5,090	57.0%	4,920	17,500
NE	9,030	77.1%	9,430	68.9%	9,230	63.9%	8,810	60.1%	8,710	57.0%	8,720	29,460
NV	13,780	77.1%	12,800	68.9%	12,230	63.9%	12,020	60.1%	11,370	57.0%	11,030	39,100
NH	8,350	77.1%	8,210	68.9%	7,810	63.9%	7,630	60.1%	7,030	57.0%	7,140	24,880
NJ	50,650	77.1%	49,080	68.9%	48,100	63.9%	47,920	60.1%	45,900	57.0%	49,370	157,330

STATE	2012	1 Year	2011	2 Year	2010	3 Year	2009	4 Year	2008	5 Year	2007	Estimated
	Estimated	Survival	Estimated	Survival	Estimated	Survival	Estimated	Survival	Estimated	Survival	Estimated	1-5 Year
	Cases		Cases		Cases		Cases		Cases		Cases	Survivors
NM	9,640	77.1%	9,630	68.9%	9,210	63.9%	8,830	60.1%	8,260	57.0%	8,030	28,950
NY	109,440	77.1%	107,260	68.9%	103,340	63.9%	101,550	60.1%	97,130	57.0%	100,960	334,710
NC	51,860	77.1%	48,870	68.9%	45,120	63.9%	42,270	60.1%	40,420	57.0%	38,210	141,850
ND	3,510	77.1%	3,560	68.9%	3,300	63.9%	3,200	60.1%	3,090	57.0%	3,340	10,820
OH	66,560	77.1%	65,060	68.9%	64,450	63.9%	62,420	60.1%	56,840	57.0%	59,220	202,370
OK	19,210	77.1%	18,980	68.9%	18,670	63.9%	18,110	60.1%	17,860	57.0%	17,170	59,590
OR	21,370	77.1%	21,180	68.9%	20,750	63.9%	19,210	60.1%	19,230	57.0%	18,630	65,080
PA	78,340	77.1%	78,030	68.9%	75,260	63.9%	74,170	60.1%	70,110	57.0%	75,130	244,370
RI	6,310	77.1%	6,090	68.9%	5,970	63.9%	6,250	60.1%	6,120	57.0%	6,360	20,110
SC	26,570	77.1%	25,510	68.9%	23,240	63.9%	22,100	60.1%	20,740	57.0%	21,370	74,450
SD	4,430	77.1%	4,430	68.9%	4,220	63.9%	4,120	60.1%	4,080	57.0%	3,990	13,680
TN	35,610	77.1%	34,750	68.9%	33,070	63.9%	32,570	60.1%	29,390	57.0%	28,440	104,260
TX	110,470	77.1%	105,000	68.9%	101,120	63.9%	98,200	60.1%	96,320	57.0%	91,020	323,150
UT	10,620	77.1%	10,530	68.9%	9,970	63.9%	8,880	60.1%	7,760	57.0%	7,660	29,690
VT	4,060	77.1%	3,950	68.9%	3,720	63.9%	3,550	60.1%	3,530	57.0%	3,500	11,990
VA	41,380	77.1%	38,720	68.9%	36,410	63.9%	34,150	60.1%	35,590	57.0%	35,090	118,150
WA	35,790	77.1%	35,360	68.9%	34,500	63.9%	32,290	60.1%	32,380	57.0%	31,080	108,840
WV	11,610	77.1%	11,080	68.9%	10,610	63.9%	10,230	60.1%	10,250	57.0%	10,490	34,530
WI	31,920	77.1%	30,530	68.9%	29,610	63.9%	27,560	60.1%	27,590	57.0%	28,130	94,170
WY	2,650	77.1%	2,680	68.9%	2,540	63.9%	2,500	60.1%	2,570	57.0%	2,340	8,290

Survival is assumed to be similar across states and is based on observed survival among cases diagnosed in SEER 17 areas during 1999-2007, followed through December 2008. The Estimated 1-5 Year Survivors do not include 2011 cases because survival is defined here as those surviving at least 1 year; the number of 1-5 year survivors is computed by the summation of 5 products: 2006 estimated cases multiplied by the 5-year observed survival rate, 2007 estimated cases multiplied by 4 year survival rate, 2008 estimated cases multiplied by 3 year survival rate, 2009 estimated cases multiplied by 2 year survival rate, and 2010 estimated cases multiplied by the 1 year survival rate.