

## Estimated 1-5 Year Cancer Survivors by State, 2013

STATE	2013	2012	1 Year	2011	2 Year	2010	3 Year	2009	4 Year	2008	5 Year	Estimated
	Estimate	Estimate	Survival	Estimate	Survival	Estimate	Survival	Estimate	Survival	Estimate	Survival	1-5 Year
	Cases	Cases		Cases		Cases		Cases		Cases		Survivors
AL	27,080	26,440	77.4%	25,530	69.2%	23,640	64.2%	24,090	60.4%	22,340	57.2%	80,640
AK	3,290	3,640	77.4%	3,090	69.2%	2,860	64.2%	2,530	60.4%	2,650	57.2%	9,840
AZ	34,010	31,990	77.4%	31,550	69.2%	29,780	64.2%	27,600	60.4%	25,540	57.2%	96,990
AR	16,330	16,120	77.4%	16,070	69.2%	15,320	64.2%	14,800	60.4%	14,840	57.2%	50,860
CA	171,330	165,810	77.4%	163,480	69.2%	157,320	64.2%	152,170	60.4%	156,530	57.2%	523,910
CO	23,410	22,820	77.4%	22,390	69.2%	21,340	64.2%	20,340	60.4%	18,900	57.2%	69,950
CT	21,180	21,530	77.4%	21,440	69.2%	20,750	64.2%	20,650	60.4%	19,190	57.2%	68,270
DE	5,370	5,340	77.4%	5,130	69.2%	4,890	64.2%	4,690	60.4%	4,590	57.2%	16,280
DC	2,920	2,980	77.4%	2,830	69.2%	2,760	64.2%	2,600	60.4%	2,560	57.2%	9,070
FL	118,320	117,580	77.4%	113,400	69.2%	107,000	64.2%	102,210	60.4%	101,920	57.2%	358,210
GA	49,280	48,130	77.4%	44,580	69.2%	40,480	64.2%	39,080	60.4%	36,980	57.2%	138,850
HI	6,650	6,610	77.4%	6,710	69.2%	6,670	64.2%	6,400	60.4%	6,310	57.2%	21,520
ID	7,670	7,720	77.4%	7,520	69.2%	7,220	64.2%	6,800	60.4%	6,430	57.2%	23,600
IL	66,090	65,750	77.4%	65,610	69.2%	63,890	64.2%	60,960	60.4%	59,130	57.2%	207,950
IN	35,550	35,060	77.4%	34,050	69.2%	33,020	64.2%	31,320	60.4%	29,550	57.2%	107,720
IA	17,480	17,010	77.4%	17,500	69.2%	17,260	64.2%	16,740	60.4%	16,150	57.2%	55,710
KS	14,370	14,090	77.4%	14,070	69.2%	13,550	64.2%	13,080	60.4%	12,520	57.2%	44,400
KY	25,100	25,160	77.4%	25,010	69.2%	24,240	64.2%	24,060	60.4%	23,270	57.2%	80,190
LA	24,930	23,480	77.4%	22,780	69.2%	20,950	64.2%	22,170	60.4%	23,360	57.2%	74,140
ME	9,190	8,990	77.4%	8,820	69.2%	8,650	64.2%	9,000	60.4%	8,140	57.2%	28,710
MD	30,680	31,000	77.4%	28,890	69.2%	27,700	64.2%	26,650	60.4%	27,380	57.2%	93,530
MA	38,250	38,470	77.4%	37,470	69.2%	36,040	64.2%	36,080	60.4%	34,470	57.2%	120,350
MI	57,560	57,790	77.4%	57,010	69.2%	55,660	64.2%	53,550	60.4%	51,150	57.2%	181,520
MN	28,410	28,060	77.4%	27,600	69.2%	25,080	64.2%	23,670	60.4%	23,160	57.2%	84,460
MS	15,830	15,190	77.4%	14,990	69.2%	14,330	64.2%	14,150	60.4%	13,400	57.2%	47,540
MO	33,950	33,440	77.4%	32,740	69.2%	31,160	64.2%	30,090	60.4%	29,390	57.2%	103,530
MT	5,450	5,550	77.4%	5,690	69.2%	5,570	64.2%	5,340	60.4%	5,090	57.2%	17,950
NE	9,060	9,030	77.4%	9,430	69.2%	9,230	64.2%	8,810	60.4%	8,710	57.2%	29,740
NV	13,830	13,780	77.4%	12,800	69.2%	12,230	64.2%	12,020	60.4%	11,370	57.2%	41,140
NH	8,470	8,350	77.4%	8,210	69.2%	7,810	64.2%	7,630	60.4%	7,030	57.2%	25,790
NJ	49,440	50,650	77.4%	49,080	69.2%	48,100	64.2%	47,920	60.4%	45,900	57.2%	159,250

STATE	2013	2012	1 Year	2011	2 Year	2010	3 Year	2009	4 Year	2008	5 Year	Estimated
	Estimate	Estimate	Survival	Estimate	Survival	Estimate	Survival	Estimate	Survival	Estimate	Survival	1-5 Year
	Cases	Cases		Cases		Cases		Cases		Cases		Survivors
NM	10,090	9,640	77.4%	9,630	69.2%	9,210	64.2%	8,830	60.4%	8,260	57.2%	30,100
NY	108,760	109,440	77.4%	107,260	69.2%	103,340	64.2%	101,550	60.4%	97,130	57.2%	342,170
NC	53,200	51,860	77.4%	48,870	69.2%	45,120	64.2%	42,270	60.4%	40,420	57.2%	151,580
ND	3,510	3,510	77.4%	3,560	69.2%	3,300	64.2%	3,200	60.4%	3,090	57.2%	11,000
OH	66,610	66,560	77.4%	65,060	69.2%	64,450	64.2%	62,420	60.4%	56,840	57.2%	208,130
OK	20,160	19,210	77.4%	18,980	69.2%	18,670	64.2%	18,110	60.4%	17,860	57.2%	61,140
OR	21,720	21,370	77.4%	21,180	69.2%	20,750	64.2%	19,210	60.4%	19,230	57.2%	67,120
PA	79,560	78,340	77.4%	78,030	69.2%	75,260	64.2%	74,170	60.4%	70,110	57.2%	247,850
RI	6,280	6,310	77.4%	6,090	69.2%	5,970	64.2%	6,250	60.4%	6,120	57.2%	20,210
SC	27,620	26,570	77.4%	25,510	69.2%	23,240	64.2%	22,100	60.4%	20,740	57.2%	78,350
SD	4,570	4,430	77.4%	4,430	69.2%	4,220	64.2%	4,120	60.4%	4,080	57.2%	14,030
TN	36,580	35,610	77.4%	34,750	69.2%	33,070	64.2%	32,570	60.4%	29,390	57.2%	109,320
TX	112,230	110,470	77.4%	105,000	69.2%	101,120	64.2%	98,200	60.4%	96,320	57.2%	337,490
UT	10,810	10,620	77.4%	10,530	69.2%	9,970	64.2%	8,880	60.4%	7,760	57.2%	31,710
VT	4,200	4,060	77.4%	3,950	69.2%	3,720	64.2%	3,550	60.4%	3,530	57.2%	12,430
VA	40,870	41,380	77.4%	38,720	69.2%	36,410	64.2%	34,150	60.4%	35,590	57.2%	123,180
WA	37,290	35,790	77.4%	35,360	69.2%	34,500	64.2%	32,290	60.4%	32,380	57.2%	112,340
WV	11,450	11,610	77.4%	11,080	69.2%	10,610	64.2%	10,230	60.4%	10,250	57.2%	35,510
WI	31,590	31,920	77.4%	30,530	69.2%	29,610	64.2%	27,560	60.4%	27,590	57.2%	97,270
WY	2,700	2,650	77.4%	2,680	69.2%	2,540	64.2%	2,500	60.4%	2,570	57.2%	8,520

Survival is assumed to be similar across states and is based on observed survival among cases diagnosed in SEER 18 areas during 2002-2008, followed through Dec 2009. The estimated 1-5 Year survivors do not include 2013 cases because survival is defined here as those surviving at least 1 year. The number of 1-5 year survivors is estimated by the summation of 5 products: 2008 estimated cases multiplied by the 5-year survival rate, 2009 estimated cases multiplied by the 4-year survival rate, 2010 estimated cases multiplied by the 3-year survival rate, 2011 estimated cases multiplied by the 2-year survival rate, and 2012 estimated cases multiplied by the 1-year survival rate.