Cancer Incidence and Mortality Rates* by Site, Race, and Ethnicity, US, 2001-2005

| Incidence | White | African American | Asian American and Pacific Islander | American Indian and Alaska Native⁺ | Hispanic/ Latino ^{‡§} |
|-----------------------|-------|---------------------|--|---------------------------------------|-----------------------------------|
| All sites | | | | | |
| Male | 551.4 | 651.5 | 354.0 | 336.6 | 419.4 |
| Female | 423.6 | 398.9 | 287.8 | 296.4 | 317.8 |
| Breast (female) | 130.6 | 117.5 | 89.6 | 75.0 | 90.1 |
| . , | 10010 | | | , | 50.1 |
| Colon & rectum | 50.0 | 71.2 | 10.0 | 16.0 | 47.2 |
| Male | 58.9 | 71.2 | 48.0 | 46.0 | 47.3 |
| Female | 43.2 | 54.5 | 35.4 | 41.2 | 32.8 |
| Kidney & renal pelvis | | | | | |
| Male | 18.8 | 21.3 | 9.1 | 19.5 | 17.4 |
| Female | 9.5 | 10.1 | 4.6 | 12.7 | 9.6 |
| Liver & bile duct | | | | | |
| Male | 8.2 | 13.2 | 21.7 | 14.4 | 15.0 |
| Female | 2.9 | 4.0 | 8.3 | 6.3 | 5.8 |
| | 2.5 | | 0.5 | 0.5 | 5.0 |
| Lung & bronchus | | | | | |
| Male | 79.3 | 107.6 | 53.9 | 54.3 | 44.2 |
| Female | 54.9 | 54.6 | 28.0 | 39.7 | 25.4 |
| Prostate | 156.7 | 248.5 | 93.8 | 73.3 | 138.0 |
| Stomach | | | | | |
| Male | 10.0 | 17.4 | 18.6 | 16.8 | 15.5 |
| Female | 4.7 | 8.9 | 10.5 | 7.7 | 9.5 |
| Uterine cervix | 8.2 | 10.8 | 8.0 | 6.9 | 13.2 |
| | | African | Asian American | American Indian | Hispanic |
| Mortality | White | American | and Pacific Islander | and Alaska Native [†] | Latino ^{‡1} |
| All sites | | | | | |
| | 220.7 | 212.0 | 120.0 | 100.0 | 150.0 |
| Male | 230.7 | 313.0 | 138.8 | 190.0 | 159.0 |
| Female | 159.2 | 186.7 | 95.9 | 142.0 | 105.2 |
| Breast (female) | 24.4 | 33.5 | 12.6 | 17.1 | 15.8 |
| Colon & rectum | | | | | |
| Male | 22.1 | 31.8 | 14.4 | 20.5 | 16.5 |
| Female | 15.3 | 22.4 | 10.2 | 14.2 | 10.8 |
| Kidney & renal pelvis | | | | | |
| Male | 6.2 | 6.1 | 2.4 | 9.3 | 5.3 |
| | 0.2 | 0.1 | 2.7 | 5.5 | 5.5 |

* Per 100,000, age adjusted to the 2000 US standard population. † Data based on Contract Health Service Delivery Areas (CHSDA), 624 counties comprising 54% of the US American Indian/Alaska Native population. ‡ Persons of Hispanic/Latino origin may be of any race. § Data unavailable from the Alaska Native Registry and Kentucky. ¶ Data unavailable from Minnesota, New Hampshire, and North Dakota.

1.2

15.2

6.6

37.5

18.5

11.0

10.1

5.9

2.2

Source: Ries LAG, Melbert D, Krapcho M, et al. (eds). SEER Cancer Statistics Review, 1975-2005, National Cancer Institute, Bethesda, MD, seer.cancer.gov/csr/1975_2005/, 2008.

2.7

10.3

3.9

93.1

39.9

59.4

11.5

5.5

4.7

Female

Female

Male Female

Prostate

Stomach Male

Female

Uterine cervix

Liver & bile duct Male

Lung & bronchus

2.8

6.7

2.9

71.3

42.0

24.6

5.0

2.5

2.3

American Cancer Society, Surveillance and Health Policy Research, 2009

4.3

10.6

6.6

50.2

33.8

21.1

9.9

5.2

3.7

2.4

11.1

5.1

35.1

14.6

20.6

8.7

4.9

3.2