

Estimated 1-5 Year Cancer Survivors 2010 (Relative Survival Rates)

STATE	2010		1 Year		2009		2 Year		2008		3 Year		2007		4 Year		2006		5 Year		2005		Estimated		
	Estimate	Cases	Survival	Estimate	Estimate	Cases	Survival	Survival	Estimate	Cases	Survival	Survival	Estimate	Cases	Survival	Survival	Estimate	Cases	Survival	Survival	Estimate	Cases	Survival	Survival	1-5 Year
AL	23,640	23,640	78.9%	24,090	22,340	69.4%	20,590	67.4%	24,390	66.1%	24,390	66.1%	24,390	66.1%	24,320	82,010									
AK	2,860	2,860	78.9%	2,530	2,650	69.4%	2,500	67.4%	2,010	66.1%	2,010	66.1%	2,010	66.1%	1,930	8,280									
AZ	29,780	29,780	78.9%	27,600	25,540	69.4%	26,270	67.4%	25,450	66.1%	25,450	66.1%	25,450	66.1%	23,880	91,460									
AR	15,320	15,320	78.9%	14,800	14,840	69.4%	14,130	67.4%	15,220	66.1%	15,220	66.1%	15,220	66.1%	14,950	52,380									
CA	157,320	157,320	78.9%	152,170	156,530	69.4%	151,250	67.4%	138,680	66.1%	138,680	66.1%	138,680	66.1%	135,030	521,240									
CO	21,340	21,340	78.9%	20,340	18,900	69.4%	19,190	67.4%	16,850	66.1%	16,850	66.1%	16,850	66.1%	16,080	65,050									
CT	20,750	20,750	78.9%	20,650	19,190	69.4%	19,780	67.4%	17,320	66.1%	17,320	66.1%	17,320	66.1%	16,920	66,790									
DE	4,890	4,890	78.9%	4,690	4,590	69.4%	4,530	67.4%	4,190	66.1%	4,190	66.1%	4,190	66.1%	3,800	15,510									
DC	2,760	2,760	78.9%	2,600	2,560	69.4%	2,540	67.4%	2,680	66.1%	2,680	66.1%	2,680	66.1%	2,820	9,340									
FL	107,000	107,000	78.9%	102,210	101,920	69.4%	106,560	67.4%	98,960	66.1%	98,960	66.1%	98,960	66.1%	96,200	358,780									
GA	40,480	40,480	78.9%	39,080	36,980	69.4%	35,440	67.4%	36,650	66.1%	36,650	66.1%	36,650	66.1%	35,650	130,510									
HI	6,670	6,670	78.9%	6,400	6,310	69.4%	6,020	67.4%	5,600	66.1%	5,600	66.1%	5,600	66.1%	4,790	20,740									
ID	7,220	7,220	78.9%	6,800	6,430	69.4%	6,140	67.4%	5,900	66.1%	5,900	66.1%	5,900	66.1%	5,490	21,890									
IL	63,890	63,890	78.9%	60,960	59,130	69.4%	62,010	67.4%	60,220	66.1%	60,220	66.1%	60,220	66.1%	59,730	214,070									
IN	33,020	33,020	78.9%	31,320	29,550	69.4%	30,040	67.4%	32,710	66.1%	32,710	66.1%	32,710	66.1%	31,900	110,120									
IA	17,260	17,260	78.9%	16,740	16,150	69.4%	16,540	67.4%	16,310	66.1%	16,310	66.1%	16,310	66.1%	15,910	57,900									
KS	13,550	13,550	78.9%	13,080	12,520	69.4%	12,760	67.4%	13,210	66.1%	13,210	66.1%	13,210	66.1%	12,930	45,700									
KY	24,240	24,240	78.9%	24,060	23,270	69.4%	22,850	67.4%	23,690	66.1%	23,690	66.1%	23,690	66.1%	23,020	82,900									
LA	20,950	20,950	78.9%	22,170	23,360	69.4%	22,540	67.4%	23,890	66.1%	23,890	66.1%	23,890	66.1%	23,280	81,560									
ME	8,650	8,650	78.9%	9,000	8,140	69.4%	8,340	67.4%	7,910	66.1%	7,910	66.1%	7,910	66.1%	7,750	29,240									
MD	27,700	27,700	78.9%	26,650	27,380	69.4%	26,390	67.4%	25,870	66.1%	25,870	66.1%	25,870	66.1%	25,450	93,450									
MA	36,040	36,040	78.9%	36,080	34,470	69.4%	34,920	67.4%	33,310	66.1%	33,310	66.1%	33,310	66.1%	33,030	121,980									
MI	55,660	55,660	78.9%	53,550	51,150	69.4%	54,410	67.4%	48,250	66.1%	48,250	66.1%	48,250	66.1%	50,220	182,810									
MN	25,080	25,080	78.9%	23,670	23,160	69.4%	25,420	67.4%	23,520	66.1%	23,520	66.1%	23,520	66.1%	22,890	84,090									
MS	14,330	14,330	78.9%	14,150	13,400	69.4%	12,470	67.4%	15,120	66.1%	15,120	66.1%	15,120	66.1%	14,970	49,620									
MO	31,160	31,160	78.9%	30,090	29,390	69.4%	29,930	67.4%	31,050	66.1%	31,050	66.1%	31,050	66.1%	30,210	106,720									
MT	5,570	5,570	78.9%	5,340	5,090	69.4%	4,920	67.4%	4,780	66.1%	4,780	66.1%	4,780	66.1%	4,910	17,790									
NE	9,230	9,230	78.9%	8,810	8,710	69.4%	8,720	67.4%	8,450	66.1%	8,450	66.1%	8,450	66.1%	8,330	30,520									
NV	12,230	12,230	78.9%	12,020	11,370	69.4%	11,030	67.4%	11,600	66.1%	11,600	66.1%	11,600	66.1%	11,120	40,550									
NH	7,810	7,810	78.9%	7,630	7,030	69.4%	7,140	67.4%	6,470	66.1%	6,470	66.1%	6,470	66.1%	6,310	24,600									
NJ	48,100	48,100	78.9%	47,920	45,900	69.4%	49,370	67.4%	43,910	66.1%	43,910	66.1%	43,910	66.1%	43,000	163,370									

STATE	2010		2009		2008		2007		2006		2005		Estimated	
	Estimate	Survival	Estimate	Survival	Estimate	Survival	Estimate	Survival	Estimate	Survival	Estimate	Survival	Estimate	Survival
	Cases		Cases		Cases		Cases		Cases		Cases		Cases	
NM	9,210	78.9%	8,830	72.5%	8,260	69.4%	8,030	67.4%	8,150	66.1%	7,780	66.1%	7,780	29,160
NY	103,340	78.9%	101,550	72.5%	97,130	69.4%	100,960	67.4%	88,230	66.1%	87,050	66.1%	87,050	337,620
NC	45,120	78.9%	42,270	72.5%	40,420	69.4%	38,210	67.4%	40,890	66.1%	40,520	66.1%	40,520	143,520
ND	3,300	78.9%	3,200	72.5%	3,090	69.4%	3,340	67.4%	3,170	66.1%	3,080	66.1%	3,080	11,260
OH	64,450	78.9%	62,420	72.5%	56,840	69.4%	59,220	67.4%	61,310	66.1%	59,680	66.1%	59,680	212,330
OK	18,670	78.9%	18,110	72.5%	17,860	69.4%	17,170	67.4%	18,640	66.1%	18,460	66.1%	18,460	63,920
OR	20,750	78.9%	19,210	72.5%	19,230	69.4%	18,630	67.4%	18,290	66.1%	17,720	66.1%	17,720	66,070
PA	75,260	78.9%	74,170	72.5%	70,110	69.4%	75,130	67.4%	73,630	66.1%	71,840	66.1%	71,840	258,600
RI	5,970	78.9%	6,250	72.5%	6,120	69.4%	6,360	67.4%	5,900	66.1%	5,870	66.1%	5,870	21,640
SC	23,240	78.9%	22,100	72.5%	20,740	69.4%	21,370	67.4%	22,530	66.1%	21,860	66.1%	21,860	76,940
SD	4,220	78.9%	4,120	72.5%	4,080	69.4%	3,990	67.4%	4,060	66.1%	3,900	66.1%	3,900	14,290
TN	33,070	78.9%	32,570	72.5%	29,390	69.4%	28,440	67.4%	32,140	66.1%	31,080	66.1%	31,080	108,950
TX	101,120	78.9%	98,200	72.5%	96,320	69.4%	91,020	67.4%	85,570	66.1%	86,880	66.1%	86,880	325,580
UT	9,970	78.9%	8,880	72.5%	7,760	69.4%	7,660	67.4%	6,620	66.1%	6,380	66.1%	6,380	26,630
VT	3,720	78.9%	3,550	72.5%	3,530	69.4%	3,500	67.4%	2,970	66.1%	3,030	66.1%	3,030	11,790
VA	36,410	78.9%	34,150	72.5%	35,590	69.4%	35,090	67.4%	34,990	66.1%	33,680	66.1%	33,680	122,950
WA	34,500	78.9%	32,290	72.5%	32,380	69.4%	31,080	67.4%	28,350	66.1%	27,350	66.1%	27,350	107,710
WV	10,610	78.9%	10,230	72.5%	10,250	69.4%	10,490	67.4%	11,280	66.1%	11,190	66.1%	11,190	37,780
WI	29,610	78.9%	27,560	72.5%	27,590	69.4%	28,130	67.4%	26,390	66.1%	26,340	66.1%	26,340	96,470
WY	2,540	78.9%	2,500	72.5%	2,570	69.4%	2,340	67.4%	2,500	66.1%	2,380	66.1%	2,380	8,720

* Survival is assumed to be similar across states and is based on cases diagnosed in SEER 17 areas during 1999-2005, followed through Dec 2006. Note: The Estimated 1-5 Year Survivors do not include 2010 cases because survival is defined here as those surviving at least 1 year; the number of 1-5 year survivors is computed by the summation of 5 products: 2005 estimated cases multiplied by 5-year survival rate, 2006 estimated cases multiplied by 4-year survival rate, 2007 estimated cases multiplied by 3 year survival rate, 2008 estimated cases multiplied by 2 year survival rate, and 2009 estimated cases multiplied by 1 year survival rate.

Estimated 1-5 Year Cancer Survivors 2010 (Observed Survival Rates)

STATE	2010		1 Year		2009		2 Year		2008		3 Year		2007		4 Year		2006		5 Year		2005		Estimated	
	Estimate	Cases	Survival	Estimate	Estimate	Cases	Survival	Estimate	Estimate	Cases	Survival	Estimate	Estimate	Cases	Survival	Estimate	Estimate	Cases	Survival	Estimate	Estimate	Cases	Survival	Survivors
AL	23,640	76.6%	24,090	68.1%	22,340	63.0%	20,590	59.2%	24,390	56.0%	24,320	74,700												
AK	2,860	76.6%	2,530	68.1%	2,650	63.0%	2,500	59.2%	2,010	56.0%	1,930	7,590												
AZ	29,780	76.6%	27,600	68.1%	25,540	63.0%	26,270	59.2%	25,450	56.0%	23,880	83,520												
AR	15,320	76.6%	14,800	68.1%	14,840	63.0%	14,130	59.2%	15,220	56.0%	14,950	47,730												
CA	157,320	76.6%	152,170	68.1%	156,530	63.0%	151,250	59.2%	138,680	56.0%	135,030	476,160												
CO	21,340	76.6%	20,340	68.1%	18,900	63.0%	19,190	59.2%	16,850	56.0%	16,080	59,520												
CT	20,750	76.6%	20,650	68.1%	19,190	63.0%	19,780	59.2%	17,320	56.0%	16,920	61,080												
DE	4,890	76.6%	4,690	68.1%	4,590	63.0%	4,530	59.2%	4,190	56.0%	3,800	14,180												
DC	2,760	76.6%	2,600	68.1%	2,560	63.0%	2,540	59.2%	2,680	56.0%	2,820	8,500												
FL	107,000	76.6%	102,210	68.1%	101,920	63.0%	106,560	59.2%	98,960	56.0%	96,200	327,290												
GA	40,480	76.6%	39,080	68.1%	36,980	63.0%	35,440	59.2%	36,650	56.0%	35,650	119,110												
HI	6,670	76.6%	6,400	68.1%	6,310	63.0%	6,020	59.2%	5,600	56.0%	4,790	18,990												
ID	7,220	76.6%	6,800	68.1%	6,430	63.0%	6,140	59.2%	5,900	56.0%	5,490	20,020												
IL	63,890	76.6%	60,960	68.1%	59,130	63.0%	62,010	59.2%	60,220	56.0%	59,730	195,130												
IN	33,020	76.6%	31,320	68.1%	29,550	63.0%	30,040	59.2%	32,710	56.0%	31,900	100,270												
IA	17,260	76.6%	16,740	68.1%	16,150	63.0%	16,540	59.2%	16,310	56.0%	15,910	52,810												
KS	13,550	76.6%	13,080	68.1%	12,520	63.0%	12,760	59.2%	13,210	56.0%	12,930	41,650												
KY	24,240	76.6%	24,060	68.1%	23,270	63.0%	22,850	59.2%	23,690	56.0%	23,020	75,590												
LA	20,950	76.6%	22,170	68.1%	23,360	63.0%	22,540	59.2%	23,890	56.0%	23,280	74,270												
ME	8,650	76.6%	9,000	68.1%	8,140	63.0%	8,340	59.2%	7,910	56.0%	7,750	26,710												
MD	27,700	76.6%	26,650	68.1%	27,380	63.0%	26,390	59.2%	25,870	56.0%	25,450	85,250												
MA	36,040	76.6%	36,080	68.1%	34,470	63.0%	34,920	59.2%	33,310	56.0%	33,030	111,330												
MI	55,660	76.6%	53,550	68.1%	51,150	63.0%	54,410	59.2%	48,250	56.0%	50,220	166,820												
MN	25,080	76.6%	23,670	68.1%	23,160	63.0%	25,420	59.2%	23,520	56.0%	22,890	76,660												
MS	14,330	76.6%	14,150	68.1%	13,400	63.0%	12,470	59.2%	15,120	56.0%	14,970	45,150												
MO	31,160	76.6%	30,090	68.1%	29,390	63.0%	29,930	59.2%	31,050	56.0%	30,210	97,220												
MT	5,570	76.6%	5,340	68.1%	5,090	63.0%	4,920	59.2%	4,780	56.0%	4,910	16,240												
NE	9,230	76.6%	8,810	68.1%	8,710	63.0%	8,720	59.2%	8,450	56.0%	8,330	27,840												
NV	12,230	76.6%	12,020	68.1%	11,370	63.0%	11,030	59.2%	11,600	56.0%	11,120	36,990												
NH	7,810	76.6%	7,630	68.1%	7,030	63.0%	7,140	59.2%	6,470	56.0%	6,310	22,490												
NJ	48,100	76.6%	47,920	68.1%	45,900	63.0%	49,370	59.2%	43,910	56.0%	43,000	149,140												

STATE	2010		1 Year		2009		2 Year		2008		3 Year		2007		4 Year		2006		5 Year		2005		Estimated	
	Estimate	Cases	Survival	Estimate	Survival	Estimate	Cases	Survival	Estimate	Cases	Survival	Estimate	Cases	Survival	Estimate	Cases	Survival	Estimate	Cases	Survival	Estimate	Cases	Survival	Estimate
NM	9,210	8,830	76.6%	8,830	68.1%	8,260	63.0%	8,030	59.2%	8,150	56.0%	7,780	26,630											
NY	103,340	101,550	76.6%	101,550	68.1%	97,130	63.0%	100,960	59.2%	88,230	56.0%	87,050	308,520											
NC	45,120	42,270	76.6%	42,270	68.1%	40,420	63.0%	38,210	59.2%	40,890	56.0%	40,520	130,880											
ND	3,300	3,200	76.6%	3,200	68.1%	3,090	63.0%	3,340	59.2%	3,170	56.0%	3,080	10,260											
OH	64,450	62,420	76.6%	62,420	68.1%	56,840	63.0%	59,220	59.2%	61,310	56.0%	59,680	193,550											
OK	18,670	18,110	76.6%	18,110	68.1%	17,860	63.0%	17,170	59.2%	18,640	56.0%	18,460	58,220											
OR	20,750	19,210	76.6%	19,210	68.1%	19,230	63.0%	18,630	59.2%	18,290	56.0%	17,720	60,300											
PA	75,260	74,170	76.6%	74,170	68.1%	70,110	63.0%	75,130	59.2%	73,630	56.0%	71,840	235,710											
RI	5,970	6,250	76.6%	6,250	68.1%	6,120	63.0%	6,360	59.2%	5,900	56.0%	5,870	19,740											
SC	23,240	22,100	76.6%	22,100	68.1%	20,740	63.0%	21,370	59.2%	22,530	56.0%	21,860	70,100											
SD	4,220	4,120	76.6%	4,120	68.1%	4,080	63.0%	3,990	59.2%	4,060	56.0%	3,900	13,040											
TN	33,070	32,570	76.6%	32,570	68.1%	29,390	63.0%	28,440	59.2%	32,140	56.0%	31,080	99,310											
TX	101,120	98,200	76.6%	98,200	68.1%	96,320	63.0%	91,020	59.2%	85,570	56.0%	86,880	297,470											
UT	9,970	8,880	76.6%	8,880	68.1%	7,760	63.0%	7,660	59.2%	6,620	56.0%	6,380	24,400											
VT	3,720	3,550	76.6%	3,550	68.1%	3,530	63.0%	3,500	59.2%	2,970	56.0%	3,030	10,780											
VA	36,410	34,150	76.6%	34,150	68.1%	35,590	63.0%	35,090	59.2%	34,990	56.0%	33,680	112,080											
WA	34,500	32,290	76.6%	32,290	68.1%	32,380	63.0%	31,080	59.2%	28,350	56.0%	27,350	98,460											
WV	10,610	10,230	76.6%	10,230	68.1%	10,250	63.0%	10,490	59.2%	11,280	56.0%	11,190	34,370											
WI	29,610	27,560	76.6%	27,560	68.1%	27,590	63.0%	28,130	59.2%	26,390	56.0%	26,340	87,990											
WY	2,540	2,500	76.6%	2,500	68.1%	2,570	63.0%	2,340	59.2%	2,500	56.0%	2,380	7,950											

* Survival is assumed to be similar across states and is based on cases diagnosed in SEER 17 areas during 1999-2005, followed through Dec 2006. Note: The Estimated 1-5 Year Survivors do not include 2010 cases because survival is defined here as those surviving at least 1 year; the number of 1-5 year survivors is computed by the summation of 5 products: 2005 estimated cases multiplied by 5-year survival rate, 2006 estimated cases multiplied by 4-year survival rate, 2007 estimated cases multiplied by 3 year survival rate, 2008 estimated cases multiplied by 2 year survival rate, and 2009 estimated cases multiplied by 1 year survival rate.