

## Estimated 1-5 Year Cancer Survivors by State in 2015, Based on Observed Survival Rates

STATE	2015	2014	1 Year	2013	2 Year	2012	3 Year	2011	4 Year	2010	5 Year	Estimated 1-5 Year Survivors
	Estimated	Estimated	Survival	Estimated	Survival	Estimated	Survival	Estimated	Survival	Estimated	Survival	
	Cases	Cases		Cases		Cases		Cases		Cases		
AL	26,150	26,770	77.7%	27,080	69.9%	26,440	65.1%	25,530	61.5%	23,640	58.4%	86,450
AK	3,700	3,750	77.7%	3,290	69.9%	3,640	65.1%	3,090	61.5%	2,860	58.4%	11,150
AZ	32,440	32,830	77.7%	34,010	69.9%	31,990	65.1%	31,550	61.5%	29,780	58.4%	106,900
AR	15,830	16,520	77.7%	16,330	69.9%	16,120	65.1%	16,070	61.5%	15,320	58.4%	53,570
CA	172,090	171,730	77.7%	171,330	69.9%	165,810	65.1%	163,480	61.5%	157,320	58.4%	553,550
CO	24,540	23,810	77.7%	23,410	69.9%	22,820	65.1%	22,390	61.5%	21,340	58.4%	75,950
CT	21,970	22,070	77.7%	21,180	69.9%	21,530	65.1%	21,440	61.5%	20,750	58.4%	71,270
DE	5,280	5,320	77.7%	5,370	69.9%	5,340	65.1%	5,130	61.5%	4,890	58.4%	17,370
DC	2,800	2,840	77.7%	2,920	69.9%	2,980	65.1%	2,830	61.5%	2,760	58.4%	9,540
FL	114,040	114,560	77.7%	118,320	69.9%	117,580	65.1%	113,400	61.5%	107,000	58.4%	380,490
GA	48,070	47,390	77.7%	49,280	69.9%	48,130	65.1%	44,580	61.5%	40,480	58.4%	153,660
HI	6,730	6,640	77.7%	6,650	69.9%	6,610	65.1%	6,710	61.5%	6,670	58.4%	22,130
ID	8,080	7,990	77.7%	7,670	69.9%	7,720	65.1%	7,520	61.5%	7,220	58.4%	25,440
IL	65,460	66,840	77.7%	66,090	69.9%	65,750	65.1%	65,610	61.5%	63,890	58.4%	218,600
IN	35,620	35,560	77.7%	35,550	69.9%	35,060	65.1%	34,050	61.5%	33,020	58.4%	115,530
IA	17,140	17,630	77.7%	17,480	69.9%	17,010	65.1%	17,500	61.5%	17,260	58.4%	57,830
KS	14,440	14,630	77.7%	14,370	69.9%	14,090	65.1%	14,070	61.5%	13,550	58.4%	47,150
KY	26,490	25,770	77.7%	25,100	69.9%	25,160	65.1%	25,010	61.5%	24,240	58.4%	83,480
LA	24,100	24,300	77.7%	24,930	69.9%	23,480	65.1%	22,780	61.5%	20,950	58.4%	77,840
ME	8,810	9,270	77.7%	9,190	69.9%	8,990	65.1%	8,820	61.5%	8,650	58.4%	29,950
MD	30,050	30,680	77.7%	30,680	69.9%	31,000	65.1%	28,890	61.5%	27,700	58.4%	99,410
MA	37,790	37,940	77.7%	38,250	69.9%	38,470	65.1%	37,470	61.5%	36,040	58.4%	125,350
MI	57,420	58,610	77.7%	57,560	69.9%	57,790	65.1%	57,010	61.5%	55,660	58.4%	190,960
MN	29,730	29,340	77.7%	28,410	69.9%	28,060	65.1%	27,600	61.5%	25,080	58.4%	92,540
MS	16,260	15,740	77.7%	15,830	69.9%	15,190	65.1%	14,990	61.5%	14,330	58.4%	50,770
MO	34,680	33,890	77.7%	33,950	69.9%	33,440	65.1%	32,740	61.5%	31,160	58.4%	110,170
MT	5,950	5,850	77.7%	5,450	69.9%	5,550	65.1%	5,690	61.5%	5,570	58.4%	18,720
NE	9,540	9,550	77.7%	9,060	69.9%	9,030	65.1%	9,430	61.5%	9,230	58.4%	30,820
NV	13,640	14,450	77.7%	13,830	69.9%	13,780	65.1%	12,800	61.5%	12,230	58.4%	44,880
NH	8,090	8,450	77.7%	8,470	69.9%	8,350	65.1%	8,210	61.5%	7,810	58.4%	27,530
NJ	51,410	51,130	77.7%	49,440	69.9%	50,650	65.1%	49,080	61.5%	48,100	58.4%	165,530

**Estimated 1-5 Year Cancer Survivors by State in 2015, Based on Observed Survival Rates (Continued)**

STATE	2015	2014	1 Year	2013	2 Year	2012	3 Year	2011	4 Year	2010	5 Year	Estimated 1-5 Year Survivors
	Estimated Cases	Estimated Cases	Survival	Estimated Cases	Survival	Estimated Cases	Survival	Estimated Cases	Survival	Estimated Cases	Survival	
NM	9,970	10,210	77.7%	10,090	69.9%	9,640	65.1%	9,630	61.5%	9,210	58.4%	32,560
NY	107,840	107,200	77.7%	108,760	69.9%	109,440	65.1%	107,260	61.5%	103,340	58.4%	356,880
NC	50,420	52,550	77.7%	53,200	69.9%	51,860	65.1%	48,870	61.5%	45,120	58.4%	168,180
ND	3,840	3,730	77.7%	3,510	69.9%	3,510	65.1%	3,560	61.5%	3,300	58.4%	11,750
OH	65,010	67,000	77.7%	66,610	69.9%	66,560	65.1%	65,060	61.5%	64,450	58.4%	219,600
OK	19,280	19,830	77.7%	20,160	69.9%	19,210	65.1%	18,980	61.5%	18,670	58.4%	64,580
OR	22,410	22,530	77.7%	21,720	69.9%	21,370	65.1%	21,180	61.5%	20,750	58.4%	71,740
PA	81,540	79,920	77.7%	79,560	69.9%	78,340	65.1%	78,030	61.5%	75,260	58.4%	260,650
RI	6,040	6,370	77.7%	6,280	69.9%	6,310	65.1%	6,090	61.5%	5,970	58.4%	20,680
SC	25,550	26,390	77.7%	27,620	69.9%	26,570	65.1%	25,510	61.5%	23,240	58.4%	86,370
SD	4,520	4,490	77.7%	4,570	69.9%	4,430	65.1%	4,430	61.5%	4,220	58.4%	14,760
TN	38,300	36,570	77.7%	36,580	69.9%	35,610	65.1%	34,750	61.5%	33,070	58.4%	117,850
TX	113,630	115,730	77.7%	112,230	69.9%	110,470	65.1%	105,000	61.5%	101,120	58.4%	363,920
UT	11,050	10,780	77.7%	10,810	69.9%	10,620	65.1%	10,530	61.5%	9,970	58.4%	35,140
VT	4,020	4,130	77.7%	4,200	69.9%	4,060	65.1%	3,950	61.5%	3,720	58.4%	13,390
VA	41,170	40,970	77.7%	40,870	69.9%	41,380	65.1%	38,720	61.5%	36,410	58.4%	132,420
WA	38,180	38,230	77.7%	37,290	69.9%	35,790	65.1%	35,360	61.5%	34,500	58.4%	120,960
WV	11,730	11,700	77.7%	11,450	69.9%	11,610	65.1%	11,080	61.5%	10,610	58.4%	37,660
WI	32,700	32,480	77.7%	31,590	69.9%	31,920	65.1%	30,530	61.5%	29,610	58.4%	104,170
WY	2,860	2,890	77.7%	2,700	69.9%	2,650	65.1%	2,680	61.5%	2,540	58.4%	8,990

Survival is assumed to be consistent across states and is based on observed survival among cases diagnosed in SEER 18 areas during 2004-2010, followed through Dec 2011. The estimated 1-5 year survivors do not include 2015 cases because survival is defined here as those surviving at least 1 year. The number of 1-5 year survivors is estimated by the summation of 5 products: 2010 estimated cases multiplied by the 5-year observed survival rate, 2011 estimated cases multiplied by the 4-year observed survival rate, 2012 estimated cases multiplied by the 3-year observed survival rate, 2013 estimated cases multiplied by the 2-year observed survival rate, and 2014 estimated cases multiplied by the 1-year observed survival rate.