

Estimated 1-5 Year Cancer Survivors by State in 2016, Based on Observed Survival Rates

STATE	2016	2015	1 Year	2014	2 Year	2013	3 Year	2012	4 Year	2011	5 Year	Estimated 1-5 Year Survivors
	Estimated Cases	Estimated Cases	Survival	Estimated Cases	Survival	Estimated Cases	Survival	Estimated Cases	Survival	Estimated Cases	Survival	
AL	27,020	26,150	78.1%	26,770	70.4%	27,080	65.7%	26,440	62.0%	25,530	59.0%	88,520
AK	3,330	3,700	78.1%	3,750	70.4%	3,290	65.7%	3,640	62.0%	3,090	59.0%	11,770
AZ	32,510	32,440	78.1%	32,830	70.4%	34,010	65.7%	31,990	62.0%	31,550	59.0%	109,240
AR	16,460	15,830	78.1%	16,520	70.4%	16,330	65.7%	16,120	62.0%	16,070	59.0%	54,200
CA	173,200	172,090	78.1%	171,730	70.4%	171,330	65.7%	165,810	62.0%	163,480	59.0%	567,120
CO	24,730	24,540	78.1%	23,810	70.4%	23,410	65.7%	22,820	62.0%	22,390	59.0%	78,670
CT	21,700	21,970	78.1%	22,070	70.4%	21,180	65.7%	21,530	62.0%	21,440	59.0%	72,610
DE	5,630	5,280	78.1%	5,320	70.4%	5,370	65.7%	5,340	62.0%	5,130	59.0%	17,730
DC	2,910	2,800	78.1%	2,840	70.4%	2,920	65.7%	2,980	62.0%	2,830	59.0%	9,620
FL	121,240	114,040	78.1%	114,560	70.4%	118,320	65.7%	117,580	62.0%	113,400	59.0%	387,260
GA	48,670	48,070	78.1%	47,390	70.4%	49,280	65.7%	48,130	62.0%	44,580	59.0%	159,420
HI	6,850	6,730	78.1%	6,640	70.4%	6,650	65.7%	6,610	62.0%	6,710	59.0%	22,360
ID	8,120	8,080	78.1%	7,990	70.4%	7,670	65.7%	7,720	62.0%	7,520	59.0%	26,200
IL	65,090	65,460	78.1%	66,840	70.4%	66,090	65.7%	65,750	62.0%	65,610	59.0%	221,080
IN	35,180	35,620	78.1%	35,560	70.4%	35,550	65.7%	35,060	62.0%	34,050	59.0%	118,040
IA	17,100	17,140	78.1%	17,630	70.4%	17,480	65.7%	17,010	62.0%	17,500	59.0%	58,150
KS	14,530	14,440	78.1%	14,630	70.4%	14,370	65.7%	14,090	62.0%	14,070	59.0%	48,060
KY	25,720	26,490	78.1%	25,770	70.4%	25,100	65.7%	25,160	62.0%	25,010	59.0%	85,680
LA	25,070	24,100	78.1%	24,300	70.4%	24,930	65.7%	23,480	62.0%	22,780	59.0%	80,310
ME	9,270	8,810	78.1%	9,270	70.4%	9,190	65.7%	8,990	62.0%	8,820	59.0%	30,220
MD	30,990	30,050	78.1%	30,680	70.4%	30,680	65.7%	31,000	62.0%	28,890	59.0%	101,490
MA	37,620	37,790	78.1%	37,940	70.4%	38,250	65.7%	38,470	62.0%	37,470	59.0%	127,310
MI	56,530	57,420	78.1%	58,610	70.4%	57,560	65.7%	57,790	62.0%	57,010	59.0%	193,390
MN	29,130	29,730	78.1%	29,340	70.4%	28,410	65.7%	28,060	62.0%	27,600	59.0%	96,220
MS	16,680	16,260	78.1%	15,740	70.4%	15,830	65.7%	15,190	62.0%	14,990	59.0%	52,440
MO	34,270	34,680	78.1%	33,890	70.4%	33,950	65.7%	33,440	62.0%	32,740	59.0%	113,300
MT	6,070	5,950	78.1%	5,850	70.4%	5,450	65.7%	5,550	62.0%	5,690	59.0%	19,140
NE	9,740	9,540	78.1%	9,550	70.4%	9,060	65.7%	9,030	62.0%	9,430	59.0%	31,290
NV	14,390	13,640	78.1%	14,450	70.4%	13,830	65.7%	13,780	62.0%	12,800	59.0%	46,010
NH	8,680	8,090	78.1%	8,450	70.4%	8,470	65.7%	8,350	62.0%	8,210	59.0%	27,850
NJ	49,750	51,410	78.1%	51,130	70.4%	49,440	65.7%	50,650	62.0%	49,080	59.0%	168,990

Estimated 1-5 Year Cancer Survivors by State in 2016, Based on Observed Survival Rates (Continued)

STATE	2016	2015	1 Year	2014	2 Year	2013	3 Year	2012	4 Year	2011	5 Year	Estimated 1-5 Year Survivors
	Estimated Cases	Estimated Cases	Survival	Estimated Cases	Survival	Estimated Cases	Survival	Estimated Cases	Survival	Estimated Cases	Survival	
NM	9,750	9,970	78.1%	10,210	70.4%	10,090	65.7%	9,640	62.0%	9,630	59.0%	33,260
NY	110,280	107,840	78.1%	107,200	70.4%	108,760	65.7%	109,440	62.0%	107,260	59.0%	362,280
NC	54,450	50,420	78.1%	52,550	70.4%	53,200	65.7%	51,860	62.0%	48,870	59.0%	172,310
ND	3,930	3,840	78.1%	3,730	70.4%	3,510	65.7%	3,510	62.0%	3,560	59.0%	12,210
OH	66,020	65,010	78.1%	67,000	70.4%	66,610	65.7%	66,560	62.0%	65,060	59.0%	221,360
OK	19,650	19,280	78.1%	19,830	70.4%	20,160	65.7%	19,210	62.0%	18,980	59.0%	65,370
OR	22,510	22,410	78.1%	22,530	70.4%	21,720	65.7%	21,370	62.0%	21,180	59.0%	73,380
PA	83,560	81,540	78.1%	79,920	70.4%	79,560	65.7%	78,340	62.0%	78,030	59.0%	266,830
RI	6,190	6,040	78.1%	6,370	70.4%	6,280	65.7%	6,310	62.0%	6,090	59.0%	20,830
SC	27,980	25,550	78.1%	26,390	70.4%	27,620	65.7%	26,570	62.0%	25,510	59.0%	88,200
SD	4,690	4,520	78.1%	4,490	70.4%	4,570	65.7%	4,430	62.0%	4,430	59.0%	15,050
TN	37,650	38,300	78.1%	36,570	70.4%	36,580	65.7%	35,610	62.0%	34,750	59.0%	122,270
TX	116,690	113,630	78.1%	115,730	70.4%	112,230	65.7%	110,470	62.0%	105,000	59.0%	374,400
UT	11,030	11,050	78.1%	10,780	70.4%	10,810	65.7%	10,620	62.0%	10,530	59.0%	36,120
VT	4,050	4,020	78.1%	4,130	70.4%	4,200	65.7%	4,060	62.0%	3,950	59.0%	13,650
VA	43,190	41,170	78.1%	40,970	70.4%	40,870	65.7%	41,380	62.0%	38,720	59.0%	136,350
WA	37,770	38,180	78.1%	38,230	70.4%	37,290	65.7%	35,790	62.0%	35,360	59.0%	124,280
WV	11,770	11,730	78.1%	11,700	70.4%	11,450	65.7%	11,610	62.0%	11,080	59.0%	38,660
WI	32,970	32,700	78.1%	32,480	70.4%	31,590	65.7%	31,920	62.0%	30,530	59.0%	106,960
WY	2,920	2,860	78.1%	2,890	70.4%	2,700	65.7%	2,650	62.0%	2,680	59.0%	9,270

Survival is assumed to be consistent across states and is based on observed survival among cases diagnosed in SEER 18 areas during 2005-2011, followed through December 2012. The estimated 1-5 year survivors do not include 2016 cases because survival is defined here as those surviving at least 1 year. The number of 1-5 year survivors is estimated by the summation of 5 products: 2011 estimated cases multiplied by the 5-year observed survival rate, 2012 estimated cases multiplied by the 4-year observed survival rate, 2013 estimated cases multiplied by the 3-year observed survival rate, 2014 estimated cases multiplied by the 2-year observed survival rate, and 2015 estimated cases multiplied by the 1-year observed survival rate.