## Figure 3. Leading Sites of New Cancer Cases and Deaths – 2016 Estimates

Estimated New Cases		Estimated Deaths	
Male	Female	Male	Female
Prostate	Breast	Lung & bronchus	Lung & bronchus
180,890 (21%)	246,660 (29%)	85,920 (27%)	72,160 (26%)
Lung & bronchus	Lung & bronchus	Prostate	Breast
117,920 (14%)	106,470 (13%)	26,120 (8%)	40,450 (14%)
Colon & rectum	Colon & rectum	Colon & rectum	Colon & rectum
70,820 (8%)	63,670 (8%)	26,020 (8%)	23,170 (8%)
Urinary bladder	Uterine corpus	Pancreas	Pancreas
58,950 (7%)	60,050 (7%)	21,450 (7%)	20,330 (7%)
Melanoma of the skin	Thyroid	Liver & intrahepatic bile duct	Ovary
46,870 (6%)	49,350 (6%)	18,280 (6%)	14,240 (5%)
Non-Hodgkin lymphoma	Non-Hodgkin lymphoma	Leukemia	Uterine corpus
40,170 (5%)	32,410 (4%)	14,130 (4%)	10,470 (4%)
Kidney & renal pelvis	Melanoma of the skin	Esophagus	Leukemia
39,650 (5%)	29,510 (3%)	12,720 (4%)	10,270 (4%)
Oral cavity & pharynx	Leukemia	Urinary bladder	Liver & intrahepatic bile duct
34,780 (4%)	26,050 (3%)	11,820 (4%)	8,890 (3%)
Leukemia	Pancreas	Non-Hodgkin lymphoma	Non-Hodgkin lymphoma
34,090 (4%)	25,400 (3%)	11,520 (4%)	8,630 (3%)
Liver & intrahepatic bile duct	Kidney & renal pelvis	Brain & other nervous system	Brain & other nervous system
28,410 (3%)	23,050 (3%)	9,440 (3%)	6,610 (2%)
All sites	All sites	All sites	All sites
841,390 (100%)	843,820 (100%)	314,290 (100%)	281,400 (100%)

Estimates are rounded to the nearest 10, and cases exclude basal cell and squamous cell skin cancers and in situ carcinoma except urinary bladder.

©2016, American Cancer Society, Inc., Surveillance Research